

Discrimination between Quantile Regression Models for Bounded Data

Alla Abdul AlSattar Hamoodat ^{1,*}, Zainab Tawfiq Hamid ², Zakariya Yahya Algamal ¹

¹*Department of Statistics and Informatics, University of Mosul, Iraq*

²*Department of Operation Research and Intelligent Techniques, University of Mosul, Iraq*

Abstract Most often when we use the term ‘bounded’, we mean a response variable that retains inherent upper and lower boundaries; for instance, it is a proportion or a strictly positive for example incomes. This constraint has implications for the type of model to be used since most traditional linear models may not respect these boundaries. Parametric quantile regression with bounded data thus comes with a framework for analysis and interpretation of how the predictor of interest influences the response variable over different quantiles while constrained by the bounds of the theoretically assumed distribution. In this paper, several parametric quantile regression models are explored and their performance is investigated under several conditions. Our Monte Carlo simulation results suggest that some of these parametric quantile regression models can bring significant improvement relative to other existing models under certain conditions.

Keywords Bounded data, beta distribution, quantile regression model, Monte Carlo simulation, Survival Analysis, SCSO

DOI: 10.19139/soic-2310-5070-2133

1. Introduction

Regression analysis is the fit of a model with the aim of expressing it as a numerical vector, that is, it works to predict the value of a numerical variable using a single estimate, the mean, and sometimes the need to predict the full range or distribution of the target variable, however, statisticians have developed increasingly advanced methods of regression, and for this reason Quantitative regression techniques have been developed.

This interest is due to several natural and anthropogenic phenomena are measured as indexes, percentages, proportions, rates and ratios, which are bounded on a certain interval, usually the unit interval. ”In many real-world applications, data is limited to a specific period of time, often within the range [0,1], and the limited nature of this data presents unique modeling challenges, prompting researchers to focus on addressing these issues, as The need to develop unit interval distributions is rapidly increasing because of their applications in engineering, economics, psychology, and biology. The unit interval or distribution bounded by the interval [0,1] is important for modeling data contained in the intervals between zero and one, such as ratios, rates, and percentages. For example, in psychology percentages and proportions are useful for judgment probabilities, which are the percentage of the mind section captured by a given region. In economics, the variable or data under study is generally limited to unit intervals, for example, market share, capital structure, and percentage of income spent on non-permanent uses. It is also noted that unit distributions have attractive bathtub-like hazard rate shapes.

Quantitative regression is one of the methods that has taken wide application in the past two decades, presented by [1], and has become attractive to researchers as it provides a framework for modeling the relationship between the response variable and the covariates using the quantitative function. The ordinary least squares regression is

*Correspondence to: Zakariya Y. Algamal (Email: zakariya.algamal@uomosul.edu.iq). Department of Statistics and Informatics, University of Mosul, Nineveh, Iraq.

that the quantitative regression estimates are more robust against the extreme values in the response measurements, that is, it is considered one of the robust methods as it gives more details of the effect of the explanatory variables on the response variable, It also has a methodology for understanding the conditional distribution of the response variable by looking at the values of some common variables at different levels (quantities), thus providing users with a more complete picture [2] The simplest definition of quantitative regression is the value that divides a set of data into groups of equal size; Thus, the quantitative values define the boundaries between groups. Statistically speaking, quantities are values taken at regular intervals from the inverse of the cumulative distribution function (CDF) of a random variable. It is important to note that the usual quantitative regression is able to approximate the conditional quantities of a response variable in the unit period, by the methods of the equivalence principle.

A series of distributions with unit interval support has been studied in the trend literature on cumulative distribution function (CDF) transformations. From the component, we mention the log-Bilal [18], inverse Gaussian unit [21], Modeling Bounded Data under a Unit Birnbaum–Saunders Distribution with Applications in Medicine and Politics [24], Lindley unit [22], In order to evaluate the effect of one or more joint searches on the average coordination distribute on bounded by the unit interval, the following distributions can be used: beta rectangular [20], and log-Lindley [22]

In applied statistics, it is very common to deal with the uncertainty of a finite phenomenon. In many fields of knowledge, we often encounter variables such as the proportions of a particular property, the scores of some ability tests, and various indicators and rates, which are located on the period (0,1), as the distributions of units specified in the interval of units are applied to model the behavior of a random variable limited by commas Temporal (0,1) has found applications in areas such as health, biology, meteorology, hydrology, financial modeling and other sciences.

The beta distribution is the well-known statistical distribution for modeling data sets on the interval (0,1) and is a suitable and useful model in many areas of statistics, but in some cases its ability to model data is not sufficient to explain it.

The beta distribution has been widely used in statistical theory and practice for more than a hundred years, and the beta distribution is the most widely used for modeling unit outcomes, i.e. the time period is between (0, 1), and beta regression is useful for understanding the effect of covariates on the average response, as it can data as rates or ratios. On the other hand, Kumaraswamy argues that the beta distribution is not faithfully proportional to hydrological random variables like daily precipitation, daily stream flow, etc. also but it is much simpler to use especially in simulation studies where probability density function, cumulative distribution function and quantitative functions can be expressed In closed form [3]. Accordingly, there are alternative distributions to the beta distribution that have been defined and applied in the literature, such as:

The Johnson SB distribution, Kumaraswamy distribution, The Log-extended exponential-geometric distribution, Unit Generalized Half Normal distribution, The unit-Birnbaum-Saunders distribution, The unit-Burr-XII distribution, The unit-Chen distribution, The unit-Half-Normal -X distribution, The unit-Gompertz distribution, The unit-Weibull distribution, and The unit-Logistic distribution.

2. Methodology

2.1. The Johnson SB distribution(johnsonsb)

Havley and Schroeder (1977) first introduced a new class of distributions called Johnson SB in the forestry literature. It is widely used in forestry to represent the empirical distributions of forest tree variables to represent the empirical distributions of forest tree variables such as diameter, height, and volume, and has since been widely used. Wide in forest diameter and height distribution [4].

Density function (f_{SB}), Distribution function F_{SB} , Quantile function Q_{SB} for the The Johnson SB distribution reparametrized in terms of the τ -th quantile, $\tau \in (0; 1)$.

$$f_{SB}(y; \alpha, \beta) = \frac{\beta}{\sqrt{2\pi} y(1-y)} \exp \left\{ -\frac{1}{2} \left[\alpha + \beta \log \left(\frac{y}{1-y} \right) \right]^2 \right\} \quad \dots \quad (1)$$

The probability density function (p.d.f.) corresponding to (??) is :

$$F_{SB}(y; \alpha, \beta) = \emptyset \left[\alpha + \beta \log \emptyset^{-1}(\tau) \right] \quad , \quad y \in (0, 1) \quad \dots \quad (2)$$

$$Q_{SB}(\tau; \alpha, \beta) = \frac{\exp \left[\frac{\emptyset^{-1}(\tau) - \alpha}{\beta} \right]}{1 + \exp \left[\frac{\emptyset^{-1}(\tau) - \alpha}{\beta} \right]} \quad \dots \quad (3)$$

where $\alpha > 0$ and $\beta > 0$ are shape parameters and $y \in (0, 1)$

2.2. leeg The Log-extended exponential-geometric distribution

It is a probability distribution derived from a two-factor expanded exponential geometric (EEG) distribution with finite support [7].

$$f_L(y; \alpha, \beta) = \frac{\alpha y^{\alpha-1} (1 + \beta)^{\beta-1}}{(1 + \beta y^\alpha)^2} \quad , \quad y \in (0, 1) \quad , \alpha > 0 \quad , \beta > -1 \quad \dots \quad (7)$$

$$F_L(y; \alpha, \beta) = \frac{y^\alpha (1 + \beta)}{1 + \beta y^\alpha} \quad \dots \quad (8)$$

where α and β are the model parameters. In particular, the case $\alpha > 0$ and $\beta \in (-1, 0)$ corresponds to the exponential-geometric distribution .

In the sequel, the random variable defined by will be referred to as theLog-extended exponential-geometric (Log-extended exponential-geometric distribution). The leeg distribution presentsan advantage with respect to the beta distribution since it does not include special functions in its formulation.

$$Q_L(\tau; \alpha, \beta) = \left[\frac{\tau}{1 + \alpha(1 - \tau)} \right]^{\frac{1}{\beta}} \quad \dots \quad (9)$$

2.3. Unit Generalized Half Normal distribution (ughn)

Let the r.v. Y follow a ughn distribution with probability density function (pdf) and cumulative distribution function (cdf) [9] :

$$f_{ughn}(y; \alpha, \beta) = \sqrt{\frac{2}{\pi}} \frac{\beta}{y(-\log(y))} \left(\frac{\log(y)}{\alpha} \right)^\beta \exp \left[-\frac{1}{2} \left(\frac{\log(y)}{\alpha} \right)^2 \right] \quad \dots \quad (10)$$

The new pdf can be obtained with transformation of the $Y = e^{-y}$ rv, where Y has GHN r.v. On the other word, a rv Y is distributed unit GHN distribution on the interval (0,1) if its log transformation, $-\log y$, is distributed GHN(α, β). We denote it with ughne(α, β). For $\alpha = 1$, unit half normal distribution is obtained. The corresponding CDF is given by :

$$F_{ughn}(y; \alpha, \beta) = 2\emptyset \left[- \left(-\frac{\log(y)}{\alpha} \right)^\beta \right] \quad , \quad y \in (0, 1) \quad \dots \quad (11)$$

respectively, where $\alpha, \beta > 0$, $\emptyset[\cdot]$ is the pdf of standard normal distribution and $\emptyset[\cdot]$ is the cdf of standard normal distribution

$$Q_{ughn}(\tau; \alpha) = \exp \left\{ -\alpha \left[-\emptyset^{-1} \left(\frac{\tau}{2} \right) \right]^{\frac{1}{\beta}} \right\} \quad \dots \quad (12)$$

2.4. The unit-Birnbaum-Saunders distribution (ubs)

Density function, distribution function, quantile function and random number generation function for the (ubs) distribution reparametrized in terms of the τ -th quantile, $\tau \in (0; 1)$ [10].

$$f_{ubs}(y; \alpha, \beta) = \frac{1}{2y\alpha\beta\sqrt{2\pi}} \left[\left(-\frac{\alpha}{\log(y)} \right)^{\frac{1}{2}} + \left(-\frac{\alpha}{\log(y)} \right)^{\frac{3}{2}} \right] \exp \left[\frac{1}{2\beta^2} \left(2 + \frac{\log(y)}{\alpha} \right) + \frac{\alpha}{\log(y)} \right] \quad \dots (17)$$

$$F_{ubs}(y; \alpha, \beta) = \frac{y^\alpha (1 + \beta)}{1 + \beta y^\alpha}, \quad 0 < x < 1 \quad \dots (13)$$

where, $\alpha > 0$ is the shape parameter

Quantile function

$$Q_{ubs}(\tau; \alpha, \beta) = 1 - \emptyset \left\{ \frac{1}{\beta} \left[-\left(\frac{\log(y)}{\alpha} \right)^{\frac{1}{2}} - \left(-\frac{\alpha}{\log(y)} \right)^{\frac{1}{2}} \right] \right\} \quad \dots (14)$$

2.5. The unit-Burr-XII distribution (uburrxii)

Let Y be a unit random variable having the UB XII distribution. The probability density function, cumulative distribution function, quantile function and random number generation function for the of Y are [11]

$$f_{uburrxii}(y; \alpha, \beta) = \frac{\alpha\beta}{y} [-\log(y)]^{\alpha-1} [1 + (-\log(y))^\alpha]^{-\beta-1} \quad \dots (15)$$

$$F_{uburrxii}(y; \alpha, \beta) = [1 + (-\log(y))^\alpha]^{-\beta} \quad \dots (16)$$

respectively, where $\alpha > 0$ and $\beta > 0$ are shape parameters. The quantile function of y Y follows by inverting (??), namely

$$Q_{uburrxii}(y; \alpha, \beta) = \exp \left[-\left(\tau^{-\frac{1}{\beta}} - 1 \right) \right]^{\frac{1}{\alpha}} \quad \dots (17)$$

Henceforth, if y is a random variable with pdf (??), we write $Y \sim \text{uburrxii}(\alpha, \beta)$. For $\alpha = 1$, the uburrxii distribution reduces to the unit Lomax distribution [21]. By taking $\beta = 1$, it is a special case of the unit log-logistic distribution [22].

2.6. The unit-Chen distribution (uchen)

This interest is due to the many natural and human phenomena being measured as indicators, percentages, proportions, rates and proportions, which are constrained to a certain period of time, usually a unit period. The need for modeling and analyzing limited data occurs in many real-life fields such as medicine, politics and psychology hence, to describe this type of data statistically, distributions with limited support are needed [12].

$$f_{uchen}(y; \alpha, \beta) = \frac{\alpha\beta}{y} (-\log(y))^{\beta-1} \exp \left\{ [-\log(y)]^\beta \exp \left\{ \alpha \left\{ 1 - \exp[-\log(y)]^\beta \right\} \right\} \right\} \quad \dots (18)$$

$$F_{uchen}(y; \alpha, \beta) = \exp \left\{ \alpha \left\{ 1 - \exp[-\log(y)]^\beta \right\} \right\}, \quad y \in (0, 1), \alpha > 0, \beta > -1 \quad \dots (19)$$

where, $\alpha > 0$ is the shape parameter and Φ is the CDF of the standard normal distribution. It is noteworthy that $\gamma = \exp(-\beta)$ is a scale parameter and is also the median of the distribution of X , since $F(\delta; \alpha, \beta) = 0.5$. In addition, the r -th moment of X is given by

$$Q_{uchen}(\tau; \alpha, \beta) = \exp \left\{ -\left[\log \left(1 - \frac{\log(\tau)}{\alpha} \right) \right]^{\frac{1}{\beta}} \right\} \quad \dots (20)$$

2.7. The unit-Half-Normal-X distribution (ughnx)

Let the r.v. Y follow a ughnx distribution with probability density function (pdf) and cumulative distribution function (cdf) :

$$f_{ughnx}(y; \alpha, \beta) = \sqrt{\frac{2}{\pi}} \frac{\beta}{y(1-y)} \left(\frac{y}{\alpha(1-y)} \right)^{\beta} \exp \left[-\frac{1}{2} \left(\frac{y}{\alpha(1-y)} \right)^{2\beta} \right] \dots (21)$$

The new pdf can be obtained with transformation of the $Y = e^{-y}$ rv, where Y has GHN r.v. On the other word, a rv Y is distributed unit GHN distribution on the interval (0,1) if its log transformation, $-\log y$, is distributed GHN(α, β). We denote it with ughne(α, β). For $\alpha = 1$, unit half normal distribution is obtained. The corresponding CDF is given by :

$$F_{ughnx}(y; \alpha, \beta) = 2\Phi \left[\frac{y}{\alpha(1-y)} \right]^{\beta} - 1, \quad y \in (0, 1) \dots (22)$$

$$Q_{ughnx}(\tau; \alpha) = \frac{\alpha [\Phi^{-1}(\frac{\tau+1}{2})]^{\frac{1}{\alpha}}}{1 + \alpha [\Phi^{-1}(\frac{\tau+1}{2})]^{\frac{1}{\alpha}}} \dots (23)$$

2.8. The unit-Gompertz distribution (ugompertz)

The Gompertz unit (UG) distribution shows a right-skewed (unimodal) density and an inverted J-shaped intensity while the hazard rate is constant and increasing and an inverted bathtub then the bathtub hazard rate. Benjamin Gompertz in 1825 introduced the Gompertz distribution in relation to human mortality and actuarial tables. Since then, it has received much attention from demographers and actuaries. This distribution is a generalization of the exponential distribution and has many real-world applications, especially in medical and actuarial studies. It has some good relationships with some well-known distributions such as the exponential, double exponential, Weibull, maximum value (Gumbel distribution) or the generalized logistic distribution.

we describe the new bounded distribution, which arises from a logarithmic transformation in the Gompertz distribution. This transformation is also considered in Grassia (1977) for the unit-Gamma distribution and Gómez-Déniz et al. (2014) for the log-Lindley distribution.

Let Y be a non negative random variable with Gompertz distribution, then its probability density function is given by [13]:

$$f_{ugompertz}(y; \alpha, \beta) = \frac{\alpha\beta}{y} \exp \{ \alpha - \beta \log(y) - \alpha \exp[-\beta \log(y)] \} \dots (24)$$

where $y \in (0, 1)$, $\beta > 0$ and $\alpha > 0$ are scale and shape parameters, respectively. for $x \in (0, 1)$. The corresponding cumulative distribution function are given by :

$$F_{ugompertz}(y; \alpha, \beta) = \exp [\alpha (1 - y^{\beta})] \dots (25)$$

The quantile function $y = Q(\tau) = F^{-1}(\tau)$, for $0 < \tau < 1$ of the ugompertz distribution is obtained by inverting Equation (F) is given by

$$Q_{ugompertz}(\tau; \alpha, \beta) = \frac{[\alpha - \log(\tau)]^{-\frac{1}{\beta}}}{\alpha} \dots (26)$$

2.9. The unit-Weibull distribution (uweibull)

we obtain the Unit-Weibull (uweibull) distribution with p.d.f :

$$f_{uweibull}(y; \alpha, \beta) = \frac{\alpha\beta}{y} [-\log(y)]^{\beta-1} \exp \left\{ -\alpha [-\log(y)]^{\beta} \right\} \dots (27)$$

and cumulative distribution function (c.d.f.) given by :

$$F_{\text{uweibull}}(y; \alpha, \beta) = \exp[\alpha(1 - y^\beta)] \quad , \quad 0 < y < 1 \quad \dots \quad (28)$$

where $\alpha > 0$ and $\beta > 0$ are shape parameters. Special cases of the uweibull distributions include. the standard uniform distribution over the interval $(0, 1)$ ($\alpha = \beta = 1$), the power function distribution ($\beta = 1$) and the unit-Rayleigh distribution ($\beta=2$). Therefore, the new distribution has connection with some well-known distributions, and hence, it can be very useful in many practical situations.

Since it is not possible to obtain a simple analytic expressions for $E(y)$, it is difficult to model the mean of Y in the absence/presence of covariates. On the other hand, the quantile function of the uweibull distribution has a simple analytic expression given by [15] :

$$Q_{\text{uweibull}}(\tau; \alpha, \beta) = \exp\left\{-\left[-\frac{\log(\tau)}{\alpha}\right]^{\frac{1}{\beta}}\right\} \quad \dots \quad (29)$$

2.10. The unit-Logistic distribution (ulogistic)

There are continuous distributions that still have limited support that needs further study, such as the L-Logistic distribution originally proposed by Tadikamalla and Johnson (1982). By transforming the logistic criterion, this construction is similar to the SB system proposed by Johnson (1949).

We say that the random variable (r.v.) y follows a ulogistic distribution, denoted by $y \sim \text{ulogistic}(\alpha, \beta)$, if its probability density function (pdf) is given by [17] :

$$f_{\text{ulogistic}}(y; \alpha, \beta) = \frac{\beta \exp(\alpha) \left(\frac{y}{1-y}\right)^{\beta-1}}{\left[1 + \exp(\alpha) \left(\frac{y}{1-y}\right)^\beta\right]^2} \quad , \quad 0 < y < 1 \quad \dots \quad (30)$$

Where , $0 < \beta < 1$ and $\alpha > 0$, Depending on the parameters α and β , the ulogistic distribution takes on a variety of shapes .

The cumulative distribution function (cdf) of the ulogistic distribution is given by :

$$F_{\text{ulogistic}}(y; \alpha, \beta) = \frac{\exp(\alpha) \left(\frac{y}{1-y}\right)^\beta}{1 + \exp(\alpha) \left(\frac{y}{1-y}\right)^\beta} \quad , \quad 0 < y < 1 \quad \dots \quad (31)$$

function The quantile function of the ulogistic distribution has a simple analytic expression given by :

$$Q_{\text{ulogistic}}(\tau; \alpha, \beta) = F^{-1}_{\text{ulogistic}}(y; \alpha, \beta) = \frac{\exp\left(-\frac{\alpha}{\beta}\right) \left(\frac{\tau}{1-\tau}\right)^{\frac{1}{\beta}}}{1 + \exp\left(-\frac{\alpha}{\beta}\right) \left(\frac{\tau}{1-\tau}\right)^{\frac{1}{\beta}}} \quad 0 < \tau < 1 \quad \dots \quad (32)$$

This would readily enable a quantile-based analysis of this model. Note that if $\beta = 1 - \beta = 0.5$, then $Q_{\text{ulogistic}}(\tau; \alpha, \beta) = \alpha$, which means that the parameter α is indeed the 50th percentile or the median of the ulogistic distribution.

3. Bayesian Information Criterion (BIC)

We obtain this formula from the formula of the AIC standard by multiplying the degree of freedom df by the amount $\text{Ln}(n)$ instead of the value 2, and the general formula of the (BIC) standard is as follows [14, 26, 27, 28] :

$$BIC = -2\text{Ln}(Lik) + \text{Ln}(n) df \quad \dots \quad (33)$$

Table 1. The best fitting model when $n=30$ for 70% >0.5 and 30% >0.5

Tau	P	Case: 70% >0.5 +30% <0.5	Case: 30% >0.5 +70% <0.5
0.2	5	Ughnx	Uchen
		Johnsonsb	Ulogistic
		Uweibull	Johnsonsb
		Ulogistic	Ugompertz
		Uburxii	Leeg
	10	Ughnx	Uchen
		Ulogistic	Ugompertz
		Leeg	Leeg
		Johnsonsb	Johnsonsb
		Ugompertz	Uweibull
0.5	5	Ughnx	Uchen
		Johnsonsb	Ulogistic
		Uweibull	Johnsonsb
		Ubs	Ugompertz
		Ulogistic	Leeg
	10	Ubs	Uchen
		Ulogistic	Ugompertz
		Johnsonsb	Johnsonsb
		Leeg	Uweibull
		Ugompertz	Ughne
0.7	5	Ughnx	Uchen
		Johnsonsb	Ulogistic
		Uweibull	Johnsonsb
		Ubs	Ugompertz
		Ulogistic	Leeg
	10	Ughne	Uchen
		Ugompertz	Ugompertz
		Ulogistic	Johnsonsb
		Johnsonsb	Uweibull
		Leeg	Ughne
0.9	5	Johnsonsb	Uchen
		Uweibull	Ulogistic
		Ubs	Johnsonsb
		Uburxii	Ugompertz
		Ugompertz	Leeg
	10	Ughne	Uchen
		Ugompertz	Ugompertz
		Ulogistic	Johnsonsb
		Johnsonsb	Uweibull
		Uweibull	Ughne

4. Simulation and Results

In this section, we explain how to apply the ideas and concepts mentioned in theoretical consideration and what are the results obtained. In order to achieve the goal of this research, we relied on Monte Carlo simulation in the R programming language to examine the performance of distributions and compare them using the Bayesian Information Criterion (BIC).

Many scenarios were applied for the simulation experiments. Different samples ($n=30,50,200$) were taken with parameters ($\tau = 0.2, 0.5, 0.7, 0.9$) and concentrations (70%, 80%, 90%) for each group.

Explanatory variables ($X = 5, 10, 15$) were generated, and the response variable Y was deduced through the model used in the simulation experiments, using regression functions in terms of the explanatory variables generated above, plus random error.

Depending on the BIC criterion, the best model are highlighted. Tables (1-9) summarize the best selected model. Several observations can be concluded.

1- For ($n=30$), regardless the value of τ , the Ughnx regression model came in the first rank for fitting the data in most cases when ($p = 5$) and when 70% of the data > 0.5 , the other hand, when 30% of the data > 0.5 , the best model is Uchen regression model. Regarding the ($p = 10$), Uchen regression model be the best.

Table 2. The best fitting model when $n=30$ for $80\% > 0.5$ and $20\% > 0.5$

Tau	P	Case: 80% >0.5 +20% <0.5	Case: 20% >0.5 +80% <0.5
0.2	5	Ughnx	Uchen
		Johnsons	Ulogistic
		Uweibull	Johnsons
		Ulogistic	Ugompertz
		Uburxii	Leeg
	10	Ughnx	Uchen
		Ulogistic	Ugompertz
		Leeg	Leeg
		Johnsons	Johnsons
		Ugompertz	Uweibull
0.5	5	Ughnx	Uchen
		Johnsons	Ulogistic
		Uweibull	Johnsons
		Ubs	Ugompertz
		Ulogistic	Leeg
	10	Ubs	Uchen
		Ulogistic	Ugompertz
		Johnsons	Johnsons
		Leeg	Uweibull
		Ugompertz	Ughne
0.7	5	Ughnx	Uchen
		Johnsons	Ulogistic
		Uweibull	Johnsons
		Ubs	Ugompertz
		Ulogistic	Leeg
	10	Ughne	Uchen
		Ugompertz	Ugompertz
		Ulogistic	Johnsons
		Johnsons	Uweibull
		Leeg	Ughne
0.9	5	Johnsons	Uchen
		Uweibull	Ulogistic
		Ubs	Johnsons
		Uburxii	Ugompertz
		Ugompertz	Leeg
	10	Ughne	Uchen
		Ugompertz	Ugompertz
		Ulogistic	Johnsons
		Johnsons	Uweibull
		Uweibull	Ughne

2- When the percentage of the data values > 0.5 is 80% , fixing ($n=30$) regardless of τ values, the best model is Uburxii regression model when ($p = 5$) . In addition , when ($p = 10$) the Uweibull regression model is the best in all cases . On the other hand , when 20% of the data values > 0.5 , the Johnsonsb regression model is the best for both ($p = 5$) and ($p = 10$)

3- When 90% of data values > 0.5 and ($n=30$) , the best models is Ughnx for the most when ($p = 5$) and ($p = 10$) , while when 10% of the data values > 0.5 , the Kum and Johnsonsb regression model is the best .

4- When n is varying , the best model is Varies regardless the percentage value of the data > 0.5 .

5- When τ is varying , in most cases , there is no effect in changing the best model .

5. Real Application

This application is based on an empirical dataset that comes from a longitudinal study on the quality of life in patients with epilepsy in Iran from March 2014 to December 2015. The Bayesian Information Criterion (BIC) are presented in Table 10. We note that the BIC value for the Ugompertz model is smaller than the BIC value for

Table 3. The best fitting model when $n=30$ for 10% >0.5 and 90% >0.5

Tau	P	Case: 90% >0.5 + 10% <0.5	Case: 10% >0.5 + 90% <0.5
0.2	5	Ughnx	Ughnx
		Johnsons b	Johnsons b
		Ulogistic	Leeg
		Uchen	Ubs
		Ughnx	Uweibull
	10	Ughnx	Kum
		Leeg	Johnsons b
		Ubs	Leeg
		Johnsons b	Uchen
		Uburxii	Ubs
0.5	5	Ughnx	Johnsons b
		Johnsons b	Leeg
		Leeg	Ubs
		Ubs	Ulogistic
		Uweibull	Uweibull
	10	Ughnx	Kum
		Ubs	Johnsons b
		Leeg	Leeg
		Uburxii	Uchen
		Johnsons b	Ubs
0.7	5	Uchen	Johnsons b
		Ughnx	Leeg
		Johnsons b	Ubs
		Ulogistic	Ulogistic
		Ubs	Uweibull
	10	Ughnx	Kum
		Ubs	Johnsons b
		Leeg	Leeg
		Uburxii	Uchen
		Johnsons b	Ubs
0.9	5	Ughnx	Johnsons b
		Johnsons b	Leeg
		Leeg	Ubs
		Ubs	Ulogistic
		Uweibull	Uweibull
	10	Ughnx	Kum
		Ubs	Johnsons b
		Leeg	Leeg
		Uburxii	Uchen
		Johnsons b	Ubs

others". So, based on the BIC criterion, the Ugompertz distribution is preferable in fitting these data than the Beta distribution for the bounded data.

6. Conclusion

This methodology of using quantile regression is thus a way of expressing the dependence of a response variable on the explanatory variable and covariates in terms of conditional quantiles. This methodology not only provides a stronger procedure to substitute the measure of the center of location of the response but also enables a better understanding of the variation of the distribution of the response as a function of the predictor values especially at different quantiles. In this paper, the author carried out a comparative analysis of the various parametric models of quantile regression under several conditions. Consequently, it was shown with the help of the Monte Carlo simulation that some of the above outlined parametric quantile regression models can provide a distinct improvement in comparison to other existing models under certain circumstance.

Table 4. The best fitting model when $n=50$ for $70\% >0.5$ and $30\% >0.5$

Tau	P	Case: 70% >0.5 + 30% <0.5	Case: 30% >0.5 + 70% <0.5
0.2	5	Ughnx	Ugompertz
		Johnsons b	Ughne
		Ugompertz	Uchen
	10	Uburrxii	Johnsons b
		Uweibull	Ulogistic
		Ugompertz	Uchen
0.5	5	Ughnx	Johnsons b
		Uweibull	Ulogistic
		Leeg	Kum
	10	Johnsons b	Leeg
		Johnsons b	Ughne
		Ugompertz	Uchen
0.7	5	Ughnx	Johnsons b
		Uweibull	Ulogistic
		Johnsons b	Kum
	10	Leeg	Leeg
		Johnsons b	Ugompertz
		Ubs	Ughne
0.9	5	Ugompertz	Uchen
		Uburrxii	Johnsons b
		Uweibull	Ulogistic
	10	Ughnx	Johnsons b
		Uweibull	Ulogistic
		Ugompertz	Kum
	5	Leeg	Leeg
		Ubs	Uchen
		Johnsons b	Ughne
	10	Ugompertz	Uchen
		Ubs	Johnsons b
		Uburrxii	Ulogistic
	5	Uweibull	Kum
		Ugompertz	Uchen
		Uweibull	Johnsons b
	10	Leeg	Ulogistic
		Ubs	Kum
		Ulogistic	Leeg

Table 5. The best fitting model when $n=50$ for $80\% > 0.5$ and $20\% > 0.5$

Tau	P	Case: $80\% > 0.5 + 20\% < 0.5$	Case: $20\% > 0.5 + 80\% < 0.5$
0.2	5	Uburrxii	Johnsons b
		Ugompertz	Kum
	10	Ughnx	Leeg
		Leeg	Ughne
		Uchen	Uchen
		Johnsons b	Johnsons b
0.5	5	Ughnx	Ugompertz
		Ugompertz	Leeg
	10	Ulogistic	Ulogistic
		Uweibull	Uchen
		Uburrxii	Johnsons b
		Ugompertz	Kum
0.7	5	Ughnx	Leeg
		Leeg	Ughne
	10	Uchen	Uchen
		Johnsons b	Johnsons b
		Ughnx	Ugompertz
		Ugompertz	Leeg
0.9	5	Ulogistic	Ulogistic
		Uweibull	Uchen
	10	Uburrxii	Johnsons b
		Ugompertz	Kum
		Ughnx	Leeg
		Leeg	Ughne
		Uchen	Uchen
		Johnsons b	Johnsons b
		Ughnx	Ugompertz
		Ugompertz	Leeg
		Ulogistic	Ulogistic
		Uweibull	Uchen

Table 6. The best fitting model when $n=50$ for $10\% > 0.5$ and $90\% > 0.5$

Tau	P	Case: 90% >0.5 +10% <0.5	Case: 10% >0.5 +90% <0.5
0.2	5	Uchen	Uburrxii
		Ubs	Kum
	10	Johnsons b	Johnsons b
		Ulogistic	Leeg
		Kum	Uchen
		Uchen	Ughnx
0.5	5	Ubs	Kum
		Johnsons b	Johnsons b
	10	Ulogistic	Ughne
		Kum	Leeg
		Uchen	Uburrxii
		Ubs	Kum
0.7	5	Johnsons b	Johnsons b
		Ulogistic	Leeg
	10	Ughnx	Uchen
		Ughnx	Ughnx
		Uweibull	Kum
		Johnsons b	Johnsons b
0.9	5	Ughne	Ughne
		Ubs	Leeg
	10	Uchen	Uburrxii
		Ubs	Kum
		Johnsons b	Johnsons b
		Ulogistic	Leeg
	5	Ughnx	Uchen
		Ughnx	Ughnx
	10	Kum	Kum
		Ubs	Johnsons b
		Uweibull	Ughne
		Johnsons b	Leeg
	5	Ubs	Uburrxii
		Johnsons b	Kum
	10	Ulogistic	Johnsons b
		Ughnx	Leeg
		Kum	Uchen
		Kum	Ughnx
	5	Ubs	Kum
		Ughnx	Johnsons b
	10	Uweibull	Ughne
		Johnsons b	Leeg

Table 7. The best fitting model when $n=200$ for $70\% > 0.5$ and $30\% > 0.5$

Tau	P	Case: 70% >0.5 + 30% <0.5	Case: 30% >0.5 + 70% <0.5
0.2	5	Uweibull	Johnsons b
		Ugompertz	Uchen
		Leeg	Kum
	10	Johnsons b	Ulogistic
		Uburxii	Ughnx
		Ughnx	Johnsons b
0.5	5	Uweibull	Ulogistic
		Johnsons b	Leeg
		Ugompertz	Uchen
	10	Ulogistic	Uweibull
		Ubs	Johnsons b
		Uweibull	Uchen
0.7	5	Ugompertz	Kum
		Leeg	Ulogistic
		Johnsons b	Ughnx
	10	Ughnx	Johnsons b
		Ubs	Ulogistic
		Uweibull	Leeg
0.9	5	Johnsons b	Uchen
		Ugompertz	Uweibull
		Ugompertz	Johnsons b
	10	Ughnx	Uchen
		Ubs	Kum
		Uweibull	Ulogistic
	5	Leeg	Ughnx
		Ughnx	Johnsons b
		Ubs	Ulogistic
	10	Uweibull	Leeg
		Johnsons b	Uchen
		Ugompertz	Uweibull
	5	Ugompertz	Johnsons b
		Ughnx	Uchen
		Ubs	Kum
	10	Uweibull	Ulogistic
		Leeg	Ughnx
		Ughnx	Johnsons b
	5	Ubs	Ulogistic
		Uweibull	Leeg
		Johnsons b	Uchen
	10	Ugompertz	Uweibull
		Ughnx	Johnsons b
		Ubs	Ulogistic

Table 8. The best fitting model when n=200 for 80% >0.5 and 20%

Tau	P	Case: 80% >0.5 +20% <0.5	Case: 20% >0.5 +80% <0.5
0.2	5	Ugompertz	Uchen
		Uweibull	Johnsons b
		Leeg	Ulogistic
	10	Uchen	Ubs
		Uburrxii	Ugompertz
		Johnsons b	Uchen
0.5	5	Uburrxii	Johnsons b
		Ulogistic	Leeg
		Kum	Ulogistic
	10	Leeg	Ubs
		Ugompertz	Uchen
		Uweibull	Johnsons b
0.7	5	Leeg	Ulogistic
		Uchen	Ubs
		Uburrxii	Ugompertz
	10	Johnsons b	Uchen
		Uburrxii	Johnsons b
		Ulogistic	Leeg
0.9	5	Kum	Ulogistic
		Leeg	Ubs
		Ugompertz	Uchen
	10	Uweibull	Johnsons b
		Leeg	Ulogistic
		Uchen	Ubs
	5	Uburrxii	Ugompertz
		Johnsons b	Uchen
		Uburrxii	Johnsons b
	10	Ulogistic	Leeg
		Kum	Ulogistic
		Leeg	Ubs

Table 9. The best fitting model when $n=200$ for $10\% > 0.5$ and 90%

Tau	P	Case: 90% >0.5 + 10% <0.5	Case: 10% >0.5 + 90% <0.5
0.2	5	Kum	Uchen
		Ughnx Johnsons b Ubs Ugompertz	Ugompertz Ulogistic Uburxii Uweibull
	10	Ughnx	Ugompertz
		Uchen Kum Ubs Johnsons b Kum	Kum Ughne Uchen Ulogistic Uchen
	5	Ughnx Johnsons b Ubs Ugompertz	Ugompertz Ulogistic Uburxii Uweibull
		Ughnx	Kum
0.5	10	Uchen Kum Ubs Johnsons b Kum	Ugompertz Ulogistic Uchen Leeg Uchen
		Ughnx Johnsons b Ubs Ugompertz	Ugompertz Ulogistic Uburxii Uweibull
	5	Ughnx	Ugompertz
		Uchen Kum Ubs Johnsons b Kum	Kum Uchen Ughne Ulogistic Uchen
	10	Ughnx Johnsons b Ubs Ugompertz	Ugompertz Ulogistic Uburxii Uweibull
		Ughnx	Ugompertz
0.7	5	Uchen Kum Ubs Johnsons b Kum	Kum Uchen Ughne Ulogistic Uchen
		Ughnx Johnsons b Ubs Ugompertz	Ugompertz Ulogistic Uburxii Uweibull
	10	Ughnx	Ugompertz
		Uchen Kum Ubs Johnsons b Kum	Kum Uchen Ughne Ulogistic Uchen
	5	Ughnx Johnsons b Ubs Ugompertz	Ugompertz Ulogistic Uburxii Uweibull
		Ughnx	Ugompertz
0.9	10	Uchen Kum Ubs Johnsons b Kum	Kum Ulogistic Uchen Ughne
		Ughnx Johnsons b Ubs Ugompertz	Ugompertz Ulogistic Uburxii Uweibull
	5	Ughnx	Ugompertz
		Uchen Kum Ubs Johnsons b	Kum Ulogistic Uchen Ughne
	10	Ughnx	Ugompertz
		Uchen Kum Ubs Johnsons b	Kum Ulogistic Uchen Ughne

Table 10. The best fitting model for the application data

Model	BIC value
Ugompertz	-1833.281
Uweibull	-1821.399
Leeg	-1813.187
Uchen	-1802.668
Uburxii	-1791.237

REFERENCES

1. Bassett Jr, Gilbert and Koenker, Roger, *Asymptotic theory of least absolute error regression*, Journal of the American Statistical Association, vol. 73, no. 363, pp. 618-622, 1978.
2. Koenker, Roger, *Quantile regression*, Cambridge university press, vol. 38, 2005.
3. Lemonte, Artur J and Moreno-Arenas, Germán, *On a heavy-tailed parametric quantile regression model for limited range response variables*, Computational Statistics, vol. 35, no. 1, pp. 379-398, 2020.
4. Kudus, Kamziah Abd and Ahmad, MI and Lapongan, Jaffirin, *Nonlinear regression approach to estimating Johnson SB parameters for diameter data*, Canadian Journal of Forest Research, vol. 29, no. 3, pp. 310-314, 1999.
5. Kumaraswamy, Ponnambalam, *A generalized probability density function for double-bounded random processes*, Journal of hydrology, vol. 46, no. 1-2, pp. 79-88, 1980.
6. Fletcher, SG and Ponnambalam, K, *Estimation of reservoir yield and storage distribution using moments analysis*, Journal of hydrology, vol. 182, no. 1-4, pp. 259-275, 1996.
7. Bassett Jr, Gilbert and Koenker, Roger, *Asymptotic theory of least absolute error regression*, Journal of the American Statistical Association, vol. 73, no. 363, pp. 618-622, 1978.
8. Korkmaz, Mustafa Çağatay, *The unit generalized half normal distribution: A new bounded distribution with inference and application*, Univ Politehnica Bucharest, 2020.
9. Cooray, Kahadawala and Ananda, Malwane MA, *A generalization of the half-normal distribution with applications to lifetime data*, JouCommunications in Statistics—Theory and Methods, vol. 37, no. 9, pp. 1323-1337, 2008.
10. Mazucheli, Josmar and Menezes, André FB and Dey, Sanku, *The unit-Birnbaum-Saunders distribution with applications*, Chilean Journal of Statistics, vol. 9, no. 1, pp. 47-57, 2018.
11. Bhatti, Fiaz Ahmad and Hamedani, Gholamhossein G and Korkmaz, Mustafa Ç and Sheng, Wenhui and Ali, Azeem, *On the Burr XII-moment exponential distribution*, Plos one, vol. 16, no. 2, pp. e0246935, 2021.
12. Korkmaz, Mustafa C and Altun, Emrah and Chesneau, Christophe and Yousof, Haitham M, *On the unit-Chen distribution with associated quantile regression and applications*, Mathematica Slovaca, vol. 72, no. 3, pp. 765-786, 2022.
13. Jafari, Ali Akbar and Tahmasebi, Saeid, *Gompertz-power series distributions*, Communications in Statistics-Theory and Methods, vol. 45, no. 13, pp. 3761-3781, 2016.
14. Li, Yingxing and Ruppert, David, *On the asymptotics of penalized splines*, Biometrika, vol. 95, no. 2, pp. 415-436, 2008.
15. Mazucheli, J and Menezes, AFB and Ghitany, ME, *The unit-Weibull distribution and associated inference*, J. Appl. Probab. Stat, vol. 13, no. 2, pp. 1-22, 2018.
16. da Paz, Rosineide F and Balakrishnan, Narayanaswamy and Bazán, Jorge Luis, *L-logistic regression models*, Brazilian Journal of Probability and Statistics, vol. 33, no. 3, pp. 455-479, 2019.
17. Bassett Jr, Gilbert and Koenker, Roger, *Asymptotic theory of least absolute error regression*, Journal of the American Statistical Association, vol. 73, no. 363, pp. 618-622, 1978.
18. Altun, E and El-Morshedy, M and Eliwa, MS, *A new regression model for bounded response variable, An alternative to the beta and unit Lindley regression models*, PLoS ONE, vol. 16, pp. 1-15, 2021.
19. Altun, E, *The log-weighted exponential regression model: Alternative to the beta regression model*, Communications in Statistics - Theory and Methods, pp. 1-16, 2019.
20. Bayes, C and Bazán, J and García, C, *A new robust regression model for proportions*, Bayesian Analysis, vol. 7, pp. 841-866, 2012.
21. Ghitany, ME and Mazucheli, J and Menezes, A and Alqallaf, F, *The unit inverse Gaussian distribution”: A new alternative to two-parameter distributions on the unit interval*, Communications in Statistics - Theory and Methods, vol. 48, pp. 3423-3438, 2019.
22. Gómez, E and Sordo, M and Calderín, E, *The log-Lindley distribution as an alternative to the beta regression model with applications in insurance*, Insurance: Mathematics and Economics, vol. 54, pp. 49-57, 2014.
23. Koenker, R and Basset, G, *Regression Quantiles*, Econometrica, vol. 46, pp. 33-50, 1978.
24. Josmar, M. and Vector, L. and Bruna, A. and Ander, F, *A New Quantile Regression for Modeling Bounded Data under a Unit Birnbaum-Saunders Distribution with Applications in Medicine and Politics*, Symmetry, vol. 13, pp. 68-75, 2021.
25. Mazucheli, J and Menezes, A and Chakraborty, S, *On the one parameter unit Lindley distribution and its associated regression model for proportion data*, Journal of Applied Statistics, vol. 46, pp. 700-714, 2019.
26. Algamil, Z.Y. and Lee, M.H. and Al-Fakih, A.M., *High-dimensional quantitative structure-activity relationship modeling of influenza neuraminidase a/PR/8/34 (H1N1) inhibitors based on a two-stage adaptive penalized rank regression*, Journal of Chemometrics, vol. 30, pp. 50-57, 2016.
27. Kahya, M. A. and Altamir, S. A. and Algamil, Z. Y., *Improving whale optimization algorithm for feature selection with a time-varying transfer function*, Numerical Algebra, Control and Optimization, vol. 11, pp. 87-98, 2020.
28. Algamil, Z. and Ali, H. M., *An efficient gene selection method for high-dimensional microarray data based on sparse logistic regression*, Electronic Journal of Applied Statistical Analysis, vol. 10, pp. 242-256, 2017.