



The Odd Generalized Rayleigh Reciprocal Weibull Family of Distributions with Applications

Regent Retrospect Musekwa *, Boikanyo Makubate, Violet Zivai Nyamajiwa

Department of Mathematics and Statistical Sciences, Botswana International University of Science and Technology, Botswana

Abstract We introduced a novel family of models in this paper, which we named the odd-generalized Rayleigh reciprocal Weibull-G (OGR-RW-G) family. This family is noteworthy because it applies the T-X model construction technique to the generalized Rayleigh reciprocal Weibull model, addressing the inflexibility limits associated with traditional models and allowing one to use any baseline distribution. We examine some valuable statistical inferences from the OGR-RW-G, including its probability density function (pdf) represented in a linear fashion, its order statistics' pdf, moments, residual life functions and Rényi entropy. Additionally, the hazard rate functions (hrfs) and pdfs of a few particular models are determined to have analytical shapes. The OGR-RW-G model parameters are determined by the widely recognized maximum likelihood estimation (MLE) technique. We also perform a simulation exercise to evaluate the performance of the MLEs. Ultimately, the utility of the OGR-RW-G family is demonstrated by using the odd generalized Rayleigh reciprocal Weibull Burr-XII (OGR-RW-BXII) example of the OGR-RW-G to three distinct datasets. In actuality, the four parameter OGR-RW-BXII outperforms the four parameter non-nested models and some nested models that are presented.

Keywords Probability weighted moment; residual life function; reverse residual life function; generalized Rayleigh Reciprocal Weibull; simulations; maximum likelihood estimation; key risk indicators.

AMS 2010 subject classifications 62E99, 60E05

DOI: 10.19139/soic-2310-5070-2194

1. Introduction

In actuarial science, probability models play a vital role in evaluating and managing financial risks related to insurance and pension programs. These models are frequently used to estimate the expected value of future claims, potential losses, and the likelihood of adverse events. They are also applicable in analyzing events such as insurance claims, fatalities, and policy cancellations. However, traditional probability distributions often fall short in accurately representing extreme events or outliers, which occur more frequently in actuarial science and financial markets. Lighter-tailed distributions, such as the normal, Weibull, and exponential distributions, may underestimate the risks associated with these extreme occurrences.

In contrast, heavy-tailed probability distributions are designed to assign greater probability to extreme values or outliers, making them better suited for modeling such phenomena. Unlike light-tailed distributions, where the probability of extreme values diminishes rapidly, heavy-tailed distributions provide a more realistic representation of data with long tails. For instance, the Pareto distribution, known for its long right tail, is widely used to model wealth and income distributions. The Student's t-distribution, with its heavier tails, is effective for hypothesis testing and handling data with outliers. The Levy distribution, which has even heavier tails than the normal distribution, finds applications in financial mathematics and physics. Similarly, the Cauchy distribution,

*Correspondence to: Regent Retrospect Musekwa (Email: mr21100062@studentmail.biust.ac.bw). Department of Mathematics, Botswana International University of Science and Technology, Palapye, Botswana

characterized by infinite variance, is often used to model phenomena with extended tails, such as network response times and earthquakes.

These heavy-tailed distributions are indispensable in fields like actuarial science, finance, economics, physics, and engineering, as they provide a more accurate reflection of real-world scenarios where extreme events occur more frequently than predicted by lighter-tailed models. Over the past few decades, significant research has been devoted to the development and enhancement of heavy-tailed distributions. Many of these distributions are constructed by introducing additional parameters to a parent cumulative density function (cdf), creating new families of distributions with thicker and longer tails. Researchers have also worked on modifying classical distributions such as the normal, Weibull, and exponential models to enhance their analytical flexibility and better capture the characteristics of heavy-tailed data [1, 2, 3, 7, 10, 11, 12, 15, 20, 23, 22].

The Odd Generalized Rayleigh Reciprocal Weibull-G (OGR-RW-G), a unique probability-based reciprocal Weibull distribution, constructed via the T-X technique [3], is offered in order to provide an adequate explanation of risk exposure under the reinsurance revenues data set. The cdf and probability distribution function (pdf) of the Generalized Rayleigh Reciprocal Weibull (GR-RW) are given by

$$F_{GR-RW}(u) = 1 - \exp \left\{ - \left[\frac{e^{-2\delta_1 u^{-\delta_2}}}{(1 - e^{-2\delta_1 u^{-\delta_2}})^2} \right] \right\},$$

and

$$\begin{aligned} f_{GR-RW}(u) &= \frac{2\delta_1 \delta_2 u^{-(\delta_2-1)} \exp \left\{ - \left[\frac{e^{-2\delta_1 u^{-\delta_2}}}{(1 - e^{-2\delta_1 u^{-\delta_2}})^2} \right] \right\}}{(1 - e^{-2\delta_1 u^{-\delta_2}})^3} \\ &\times [1 - e^{-2\delta_1 u^{-\delta_2}} (1 - e^{-2\delta_1 u^{-\delta_2}})], \end{aligned}$$

respectively, for $u > 0$, $\delta_1 > 0$, and $\delta_2 > 0$.

Now, the cdf and pdf of the Odd Generalized Rayleigh Reciprocal Weibull-G (OGR-RW-G) distribution are given by

$$\begin{aligned} F_{OGR-RW-G}(y) &= \int_0^{\nabla} f_{GR-RW}(u) du \\ &= 1 - \exp \left\{ - \left[\frac{e^{-2\delta_1 \nabla^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla^{-\delta_2}})^2} \right] \right\}, \end{aligned} \quad (1)$$

and

$$\begin{aligned} f_{OGR-RW-G}(y) &= \frac{2\delta_1 \delta_2 \nabla^{-(\delta_2-1)} \nabla' \exp \left\{ - \left[\frac{e^{-2\delta_1 \nabla^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla^{-\delta_2}})^2} \right] \right\}}{(1 - e^{-2\delta_1 \nabla^{-\delta_2}})^3} \\ &\times [1 - e^{-2\delta_1 \nabla^{-\delta_2}} (1 - e^{-2\delta_1 \nabla^{-\delta_2}})], \end{aligned} \quad (2)$$

respectively, where δ_1, δ_2 are non-negative scale and shape parameters respectively, $y \in \mathbb{R}$, $\nabla = \frac{G(y; \underline{\omega})}{\bar{G}(y; \underline{\omega})}$, $\nabla' = \frac{g(y; \underline{\omega})}{G^2(y; \underline{\omega})}$, $G(y; \underline{\omega})$ is a baseline cdf, $\bar{G}(y; \underline{\omega})$ is a survival function and $\underline{\omega}$ is a vector of parameters. The corresponding hazard rate function (hrf) is given by

$$\begin{aligned} h(y) &= \frac{2\delta_1 \delta_2 \nabla^{-(\delta_2-1)} g(y; \underline{\omega}) \exp \left\{ - \left[\frac{e^{-2\delta_1 \nabla^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla^{-\delta_2}})^2} \right] \right\}}{\bar{G}^2(y; \underline{\omega}) (1 - e^{-2\delta_1 \nabla^{-\delta_2}})^3} \\ &\times \exp \left\{ \left[\frac{e^{-2\delta_1 \nabla^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla^{-\delta_2}})^2} \right] \right\} [1 - e^{-2\delta_1 \nabla^{-\delta_2}} (1 - e^{-2\delta_1 \nabla^{-\delta_2}})]. \end{aligned}$$

Using the generalized binomial expansion and the power series, the pdf in equation (2) can be expressed as

$$f_{OGR-RW-G}(y) = \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* g_{\tau}^*(y; \underline{\omega}), \quad (3)$$

where

$$\begin{aligned} \chi_{\tau}^* &= \sum_{d,e=0}^{\infty} \frac{(-1)^{a+b+c+d+e} [2\delta_1(a+b+c+1)]^d}{(-(d\delta_2 + \delta_2) + h)a!d!} \\ &\times \binom{2a+b+2}{b} \binom{1}{c} \binom{d\delta_2 + \delta_2 + e - 4}{e}, \end{aligned} \quad (4)$$

and

$$g_{\tau}^*(y; \underline{\omega}) = \tau g(y; \underline{\omega}) G^{\tau-1}(y; \underline{\omega}), \quad (5)$$

is an exponentiated-G (Exp-G) distribution with power parameter $\tau = -(d\delta_2 + \delta_2) + h$. Similarly, the pdf of the k^{th} order statistic of the OGR-RW-G family of distributions can be written as

$$f_{k:m}(y) = \sum_{a,b,c=0}^{\infty} \phi_{\tau}^* g_{\tau}^*(y; \underline{\omega}), \quad (6)$$

where

$$\begin{aligned} \phi_{\tau}^* &= \frac{m!}{(k-1)!(m-k)!} \sum_{l,p,d,e=0}^{\infty} \frac{(-1)^{l+p+a+b+c+d+e} (p+1)^a [2\delta_1(a+b+c+1)]^d}{(-(d\delta_2 + \delta_2) + h)(p+1)^{-a} a!d!} \\ &\times \binom{m-k}{l} \binom{k+l-1}{p} \binom{2a+b+2}{b} \binom{1}{c} \binom{d\delta_2 + \delta_2 + e - 4}{e}. \end{aligned} \quad (7)$$

Thus, the pdf of the order statistic of the OGR-RW-G can be written as a linear combination of Exp-G with power parameter τ .

2. Special Cases

The OGR-RW-G family of distributions' special instances are discussed in this section. Examples of log-logistic, exponential, Fréchet and Burr-XII distributions for the baseline distribution function are given. The pdf and hrf plots are also shown.

2.1. Odd-Generalized Rayleigh Reciprocal Weibull-Log-Logistic

Suppose we have a log-logistic baseline distribution with cdf and pdf given by $G(y; \alpha) = 1 - (1 + y^{\alpha})^{-1}$ and $g(y; \alpha) = \alpha y^{\alpha-1} (1 + y^{\alpha})^{-2}$ respectively for $y, \alpha > 0$. Now, the cdf and pdf of the Odd-Generalized Rayleigh Reciprocal Weibull-Log-Logistic (OGR-RW-LLoG) distribution are given by

$$F_{OGR-RW-LLoG}(y) = 1 - \exp \left\{ - \left[\frac{e^{-2\delta_1 \nabla_1^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla_1^{-\delta_2}})^2} \right] \right\}, \quad (8)$$

and

$$\begin{aligned} f_{OGR-RW-LLoG}(y) &= \frac{2\delta_1 \delta_2 \nabla_1^{-(\delta_2+1)} \nabla_1' e^{-2\delta_1 \nabla_1^{-\delta_2}} (1 + e^{-2\delta_1 \nabla_1^{-\delta_2}})}{(1 - e^{-2\delta_1 \nabla_1^{-\delta_2}})^3} \\ &\times \exp \left\{ - \left[\frac{e^{-2\delta_1 \nabla_1^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla_1^{-\delta_2}})^2} \right] \right\}, \end{aligned} \quad (9)$$

respectively, for $\alpha, \delta_1, \delta_2 > 0$, $\nabla_1 = \frac{1-(1+y^\alpha)^{-1}}{(1+y^\alpha)^{-1}}$ and $\nabla'_1 = \frac{\alpha y^{\alpha-1}(1+y^\alpha)^{-2}}{(1+y^\alpha)^{-2}}$.




Figure 1. The pdf and hrf plots of the OGR-RW-LLoG distribution

2.2. Odd-Generalized Rayleigh Reciprocal Weibull - Exponential

By taking an exponential baseline distribution with cdf and pdf given by $G(y; \alpha) = 1 - e^{-\alpha y}$ and $g(y; \alpha) = \alpha e^{-\alpha y}$ respectively for $y, \alpha > 0$. Now, the cdf and pdf of the Odd-Generalized Rayleigh Reciprocal Weibull-Exponential (OGR-RW-E) distribution are given by

$$F_{OGR-RW-E}(y) = 1 - \exp \left\{ - \left[\frac{e^{-2\delta_1 \nabla_2^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla_2^{-\delta_2}})^2} \right] \right\}, \tag{10}$$

and

$$f_{OGR-RW-E}(y) = \frac{2\delta_1 \delta_2 \nabla_2^{-(\delta_2+1)} \nabla'_2 e^{-2\delta_1 \nabla_2^{-\delta_2}} (1 + e^{-2\delta_1 \nabla_2^{-\delta_2}})}{(1 - e^{-2\delta_1 \nabla_2^{-\delta_2}})^3} \times \exp \left\{ - \left[\frac{e^{-2\delta_1 \nabla_2^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla_2^{-\delta_2}})^2} \right] \right\}, \tag{11}$$

respectively, for $\alpha, \delta_1, \delta_2 > 0$, $\nabla_2 = \frac{1-e^{-\alpha y}}{e^{-\alpha y}}$ and $\nabla'_2 = \frac{\alpha e^{-\alpha y}}{e^{-2\alpha y}}$.

2.3. Odd-Generalized Rayleigh Reciprocal Weibull-Frétchet

Suppose our baseline distribution is Frétchet with cdf and pdf given by $G(y; \lambda, \alpha) = e^{-\lambda y^{-\alpha}}$ and $g(y; \lambda, \alpha) = \lambda \alpha y^{-(\alpha+1)} e^{-\lambda y^{-\alpha}}$ respectively for $y, \lambda, \alpha > 0$. Now, the cdf and pdf of the Odd-Generalized Rayleigh Reciprocal Weibull-Frétchet (OGR-RW-Fr) distribution are given by

$$F_{OGR-RW-Fr}(y) = 1 - \exp \left\{ - \left[\frac{e^{-2\delta_1 \nabla_3^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla_3^{-\delta_2}})^2} \right] \right\}, \tag{12}$$




Figure 2. The pdf and hrf plots of the OGR-RW-E distribution

and

$$f_{OGR-RW-Fr}(y) = \frac{2\delta_1\delta_2\nabla_3^{-(\delta_2+1)}\nabla_3'e^{-2\delta_1\nabla_3^{-\delta_2}}(1+e^{-2\delta_1\nabla_3^{-\delta_2}})}{(1-e^{-2\delta_1\nabla_3^{-\delta_2}})^3} \times \exp\left\{-\left[\frac{e^{-2\delta_1\nabla_3^{-\delta_2}}}{(1-e^{-2\delta_1\nabla_3^{-\delta_2}})^2}\right]\right\}, \tag{13}$$

respectively, for $\lambda, \alpha, \delta_1, \delta_2 > 0$, $\nabla_3 = \frac{e^{-\lambda y^{-\alpha}}}{1-e^{-\lambda y^{-\alpha}}}$ and $\nabla_3' = \frac{\lambda\alpha y^{-(\alpha+1)}e^{-\lambda y^{-\alpha}}}{(1-e^{-\lambda y^{-\alpha}})^2}$.

2.4. Odd-Generalized Rayleigh Reciprocal Weibull-Burr-XII

Suppose our baseline distribution is a Burr-XII distribution with cdf and pdf given by $G(y) = 1 - (1 + y^\beta)^{-\alpha}$ and $g(y) = \alpha\beta y^{\beta-1}(1 + y^\beta)^{-(\alpha+1)}$ respectively for $y, \alpha, \beta > 0$. Now, the cdf and pdf of the Odd-Generalized Rayleigh Reciprocal Weibull-Burr-XII (OGR-RW-BXII) distribution are given by

$$F_{OGR-RW-BXII}(y) = 1 - \exp\left\{-\left[\frac{e^{-2\delta_1\nabla_4^{-\delta_2}}}{(1-e^{-2\delta_1\nabla_4^{-\delta_2}})^2}\right]\right\}, \tag{14}$$

and

$$f_{OGR-RW-BXII}(y) = \frac{2\delta_1\delta_2\nabla_4^{-(\delta_2+1)}\nabla_4'e^{-2\delta_1\nabla_4^{-\delta_2}}(1+e^{-2\delta_1\nabla_4^{-\delta_2}})}{(1-e^{-2\delta_1\nabla_4^{-\delta_2}})^3} \times \exp\left\{-\left[\frac{e^{-2\delta_1\nabla_4^{-\delta_2}}}{(1-e^{-2\delta_1\nabla_4^{-\delta_2}})^2}\right]\right\}, \tag{15}$$

respectively, for $\alpha, \beta, \delta_1, \delta_2 > 0$, $\nabla_4 = \frac{1-(1+y^\beta)^{-\alpha}}{(1+y^\beta)^{-\alpha}}$ and $\nabla_4' = \frac{\alpha\beta y^{\beta-1}(1+y^\beta)^{-(\alpha+1)}}{(1+y^\beta)^{-2\alpha}}$.




Figure 3. The pdf and hrf plots of the OGR-RW-Fr distribution




Figure 4. The pdf and hrf plots of the OGR-RW-BXII distribution

3. Statistical Properties

Assessing the statistical properties of a model is essential for evaluating its performance, reliability, and applicability in both scientific and practical contexts. Key properties, such as measures of central tendency, variability, goodness-of-fit, and error metrics, offer valuable insights into the model’s ability to represent the underlying data or processes it is designed to simulate. These evaluations help identify potential biases, measure

prediction accuracy, and detect issues like overfitting or underfitting. Moreover, statistical analysis facilitates comparisons between models, aiding in the selection of the most suitable one for a specific task. Understanding these properties ensures that the model's outputs are interpretable, reproducible, and valid for informed decision-making or further research. In this section, we outline some of the key statistical properties.

The r^{th} moment of $Y \sim \text{OGR-RW-G}$, say μ'_r follows from equation (3) as

$$\mu'_r = E(Y^r) = \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* E(Y_{\tau}^r). \quad (16)$$

The mean and variance can be easily derived from equation (16). The moment generating function (mgf) of the OGR-RW-G distribution can be derived from equation (3) and is given by

$$M_Y(t) = \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* M_{\tau}(t), \quad (17)$$

where $M_{\tau}(t)$ is the MGF of Y_{τ} . Hence, $M_Y(t)$ can be determined from the MGF of Exp-G.

The r^{th} incomplete moment can be obtained as follows;

$$I_r(t) = \int_0^t y^r f_{\text{OGR-RW-G}}(y) dy = \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* \int_0^t y^r g_{\tau}^*(y; \underline{\omega}) dy. \quad (18)$$

The probability weighted moment (PWM) of a random variable $Y \sim \text{OGR-RW-G}$, say $(\varrho_{r,j})$ is given by

$$\begin{aligned} \varrho_{r,j} &= E(Y^r F^j(Y)) = \int_{-\infty}^{\infty} y^r F_{\text{OGR-RW-G}}^j(y; \delta, \gamma, \beta, \kappa) f_{\text{OGR-RW-G}}(y; \delta, \gamma, \beta, \kappa) dy \\ &= \sum_{a,b,c=0}^{\infty} Q_{\tau}^* \int_{-\infty}^{\infty} y^r g_{i+m+1}^*(y; \kappa) dy = \sum_{a,b,c=0}^{\infty} Q_{\tau}^* E[Y_{\tau}^r], \end{aligned} \quad (19)$$

where

$$\begin{aligned} Q_{\tau}^* &= \sum_{p,d,e=0}^{\infty} \frac{(-1)^{p+a+b+c+d+e} (p+1)^a [2\delta_1(a+b+c+1)]^d}{(-(d\delta_2 + \delta_2) + h)(p+1)^{-a} a! d!} \\ &\times \binom{j}{p} \binom{2a+b+2}{b} \binom{1}{c} \binom{d\delta_2 + \delta_2 + e - 4}{e}, \end{aligned} \quad (20)$$

$E(Y_{\tau}^r)$ is the r^{th} moment of Exponentiated-G (Exp-G) density with power parameter $\tau = -(d\delta_2 + \delta_2) + h$ and $g_{i+m+1}^*(y; \kappa)$ is given in equation (5).

If n is an integer value greater than 1 ($n > 1$) and $y > t$, then the r^{th} moment of residual life of the OGR-RW-G family of distributions is given by;

$$v_r^*(t) = \frac{1}{\bar{F}(y; \underline{\omega})} \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* \sum_{s=0}^{\infty} \binom{n}{s} (-t)^s \int_t^{\infty} y^{n-s} g_{\tau}^*(y; \underline{\omega}) dy, \quad (21)$$

where $g_{\tau}^*(y; \underline{\omega})$ is pdf of an Exp-G distribution with power parameter τ given in equation (5), $\bar{G}(y; \underline{\omega})$ is the survival function and the coefficients of χ_{τ}^* are given in equation (4).

Now, if n is an integer value greater than 1 ($n > 1$) and $y < t$, then the r^{th} moment of reverse residual life of the OGR-RW-G family of distributions is given by;

$$\theta_r^*(t) = \frac{1}{F(y; \underline{\omega})} \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* \sum_{s=0}^{\infty} \binom{n}{s} (t)^{n-s} (-1)^s \int_0^t y^s g_{\tau}^*(y; \underline{\omega}) dy, \quad (22)$$

where $g_\tau^*(y; \underline{\omega})$ is a pdf of an Exp-G distribution with power parameter (τ) given in equation (5) and the coefficients of χ_τ^* are given in equation (4).

Let $X \sim \text{OGR-RW-G}(\delta_{11}, \delta_{21}, \underline{\omega}_1)$ and $Y \sim \text{OGR-RW-G}(\delta_{12}, \delta_{22}, \underline{\omega}_2)$ respectively, the reliability model of the OGR-RW-G family of distributions is given by

$$\begin{aligned}
 R &= \int_{x=0}^{\infty} \frac{2\delta_{11}\delta_{21}\nabla_1^{-(\delta_{21}+1)}\nabla_1' e^{-2\delta_{11}\nabla_1^{-\delta_{21}}}(1 + e^{-2\delta_{11}\nabla_1^{-\delta_{21}}})}{(1 - e^{-2\delta_{11}\nabla_1^{-\delta_{21}}})^3} \\
 &\times \exp\left\{-\left[\frac{e^{-2\delta_{11}\nabla_1^{-\delta_{21}}}}{(1 - e^{-2\delta_{11}\nabla_1^{-\delta_{21}}})^2}\right]\right\} \\
 &\times \left(1 - \exp\left\{-\left[\frac{e^{-2\delta_{12}\nabla_2^{-\delta_{22}}}}{(1 - e^{-2\delta_{12}\nabla_2^{-\delta_{22}}})^2}\right]\right\}\right) dx, \tag{23}
 \end{aligned}$$

where $\nabla_1 = \frac{G(y;\underline{\omega}_1)}{G(y;\underline{\omega}_1)}$, $\nabla_1' = \frac{g(y;\underline{\omega}_1)}{G^2(y;\underline{\omega}_1)}$ and $\nabla_2 = \frac{G(y;\underline{\omega}_2)}{G(y;\underline{\omega}_2)}$.

The Rényi Entropy for the OGR-RW-G family of distribution given by;

$$\begin{aligned}
 I_R(\nu) &= \frac{1}{1-\nu} \log \left[(2\delta_1\delta_2)^\nu \sum_{a,b,c,d,e=0}^{\infty} \frac{(-1)^{a+b+c+d+e} \nu^a [2\delta_1(a+b+c+1)]^d}{a!d!} \right. \\
 &\times \binom{2a+b+3\nu-1}{b} \binom{\nu}{c} \binom{d\delta_2+\delta_2\nu-\nu+e-1}{e} \\
 &\times \left. \int_0^\infty g^\nu(x; \zeta) G^\tau(y; \underline{\omega}) dy \right], \tag{24}
 \end{aligned}$$

where $\tau = -(d\delta_2 + \delta_2) + h$, $\nu \neq 1$ and $\nu > 0$.

4. Key Risk Indicators

In this section, we analyze five Key Risk Indicators (KRIs) for our new model: Value-at-Risk (VaR), Tail-Value-at-Risk (TVaR), Conditional-Value-at-Risk (CVaR), Tail Variance (TV), and Tail Mean-Variance (TMV). These metrics are critical for quantifying and managing financial and operational risks, particularly in environments characterized by extreme events or significant uncertainties. They provide valuable insights into potential losses under adverse conditions by focusing on the tail of the distribution, where the most severe outcomes are concentrated.

These indicators not only assess the likelihood of losses but also capture their severity and variability, making them indispensable tools for stress testing, regulatory compliance, and developing effective risk mitigation strategies. By incorporating these measures, organizations can enhance their preparedness for low-probability, high-impact events and make informed decisions to ensure stability and resilience in the face of uncertainty. Let $Y \sim \text{OGR-RW-G}$ denote a loss random variable.

4.1. VAR Indicator

The VAR of Y at the 100 α % confidence level, say $\text{VAR}(Y; \alpha)$ is given by

$$\text{VAR}(Y; \alpha) = G^{-1} \left(\frac{\psi(\alpha)}{1 + \psi(\alpha)} \right), \tag{25}$$

for $\psi(\alpha) = \left[\frac{\log\left\{\frac{-\log(1-\alpha)}{1-\log(1-\alpha)}\right\}}{-2\delta_1} \right]^{-\delta_2^{-1}}$.

4.2. TVAR Indicator

The TVAR of Y at the $100\alpha\%$ confidence level, say $\text{TVAR}(Y; \alpha)$ is given by

$$\begin{aligned} \text{TVAR}(Y; \alpha) &= E[Y|Y > \text{VAR}(Y; \alpha)] = (1 - \alpha)^{-1} \int_{\text{VAR}(Y; \alpha)}^{\infty} y f_{\text{OGR-RW-G}}(y) dy \\ &= (1 - \alpha)^{-1} \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* \int_{\text{VAR}(Y; \alpha)}^{\infty} y g_{\tau}^*(y; \underline{\omega}) dy, \end{aligned} \quad (26)$$

where χ_{τ}^* is given in equation (4), $g_{\tau}^*(y; \underline{\omega})$ is given in equation (5) and $\tau = -(d\delta_2 + \delta_2) + h$ is the power parameter.

4.3. CVAR Indicator

The CVAR of Y at the $100\alpha\%$ confidence level, say $\text{CVAR}(Y; \alpha)$ is given by

$$\begin{aligned} \text{CVAR}(Y; \alpha) &= E[Y|Y < \text{VAR}(Y; \alpha)] = \frac{1}{\alpha} \int_{-\infty}^{\text{VAR}(Y; \alpha)} y f_{\text{OGR-RW-G}}(y) dy \\ &= \frac{1}{\alpha} \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* \int_{-\infty}^{\text{VAR}(Y; \alpha)} y g_{\tau}^*(y; \underline{\omega}) dy, \end{aligned} \quad (27)$$

where χ_{τ}^* is given in equation (4), $g_{\tau}^*(y; \underline{\omega})$ is given in equation (5) and $\tau = -(d\delta_2 + \delta_2) + h$ is the power parameter.

4.4. TV Risk Indicator

The TV risk indicator of Y at the $100\alpha\%$ confidence level, say $\text{TV}(Y; \alpha)$ is given by

$$\begin{aligned} \text{TV}(Y; \alpha) &= E[Y^2|Y > \text{VAR}(Y; \alpha)] - \left(\text{TVAR}(Y; \alpha) \right)^2 \\ &= (1 - \alpha)^{-1} \int_{\text{VAR}(Y; \alpha)}^{\infty} y^2 f_{\text{OGR-RW-G}}(y) dy - \left(\text{TVAR}(Y; \alpha) \right)^2 \\ &= (1 - \alpha)^{-1} \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* \int_{\text{VAR}(Y; \alpha)}^{\infty} y^2 g_{\tau}^*(y; \underline{\omega}) dy - \left(\text{TVAR}(Y; \alpha) \right)^2, \end{aligned} \quad (28)$$

where χ_{τ}^* is given in equation (4), $g_{\tau}^*(y; \underline{\omega})$ is given in equation (5) and $\tau = -(d\delta_2 + \delta_2) + h$ is the power parameter.

4.5. TMV Risk Indicator

The TMV risk indicator of Y at the $100\alpha\%$ confidence level, say $\text{TMV}(Y; \alpha)$ is given by

$$\begin{aligned} \text{TMV}(Y; \alpha) &= E[(Y - \text{TVAR}(Y; \alpha))^2|Y > \text{VAR}(Y; \alpha)] \\ &= (1 - \alpha)^{-1} \int_{\text{VAR}(Y; \alpha)}^{\infty} (y - \text{TVAR}(Y; \alpha))^2 f_{\text{OGR-RW-G}}(y) dy \\ &= (1 - \alpha)^{-1} \sum_{a,b,c=0}^{\infty} \chi_{\tau}^* \int_{\text{VAR}(Y; \alpha)}^{\infty} g_{\tau}^*(y; \underline{\omega}) (y - \text{TVAR}(Y; \alpha))^2 dy, \end{aligned} \quad (29)$$

where χ_{τ}^* is given in equation (4), $g_{\tau}^*(y; \underline{\omega})$ is given in equation (5) and $\tau = -(d\delta_2 + \delta_2) + h$ is the power parameter.

5. Maximum Likelihood Estimation

Let $X \sim \text{OGR-RW-G}(\delta_1, \delta_2, \omega)$ and $\Delta = (\delta_1, \delta_2, \omega)^T$ the vector of model parameters, then the log-likelihood function $\ell_n(\Delta) = \ell_n$ based on a random sample of size n from the OGR-RW-G family of distributions is given by

$$\begin{aligned} \ell_n &= n \log(2\delta_1\delta_2) - (\delta_2 + 1) \sum_{i=1}^n \log(\nabla) + \sum_{i=1}^n \log(\nabla') - 2 \sum_{i=1}^n \log(\delta_1 \nabla^{-\delta_2}) \\ &+ \sum_{i=1}^n \log(1 + e^{-2\delta_1 \nabla^{-\delta_2}}) - 3 \sum_{i=1}^n \log(1 - e^{-2\delta_1 \nabla^{-\delta_2}}) - \sum_{i=1}^n \left[\frac{e^{-2\delta_1 \nabla^{-\delta_2}}}{1 - e^{-2\delta_1 \nabla^{-\delta_2}}} \right]. \end{aligned}$$

The score equations for the OGR-RW-G model are

$$\begin{aligned} \frac{\partial \ell_n}{\partial \delta_1} &= \frac{n}{\delta_1} - 2 \sum_{i=1}^n \nabla^{-\delta_2} - 2 \sum_{i=1}^n \frac{\nabla^{-\delta_2} e^{-2\delta_1 \nabla^{-\delta_2}}}{1 + e^{-2\delta_1 \nabla^{-\delta_2}}} \\ &- 6 \sum_{i=1}^n \frac{\nabla^{-\delta_2} e^{-2\delta_1 \nabla^{-\delta_2}}}{1 - e^{-2\delta_1 \nabla^{-\delta_2}}} + 2 \sum_{i=1}^n \frac{\nabla^{-\delta_2} e^{-2\delta_1 \nabla^{-\delta_2}}}{(1 - e^{-2\delta_1 \nabla^{-\delta_2}})^2}, \\ \frac{\partial \ell_n}{\partial \delta_2} &= \frac{n}{\delta_2} - \sum_{i=1}^n \log(\nabla) - 2 \sum_{i=1}^n \nabla^{-\delta_2} - 2 \sum_{i=1}^n \delta_1 \nabla^{-\delta_2} \log(\nabla) \\ &- 2 \sum_{i=1}^n \frac{\delta_1 \nabla^{-\delta_2} e^{-2\delta_1 \nabla^{-\delta_2}} \log(\nabla)}{1 + e^{-2\delta_1 \nabla^{-\delta_2}}} - 6 \sum_{i=1}^n \frac{\delta_1 \nabla^{-\delta_2} e^{-2\delta_1 \nabla^{-\delta_2}} \log(\nabla)}{1 - e^{-2\delta_1 \nabla^{-\delta_2}}} \\ &+ 2 \sum_{i=1}^n \frac{\delta_1 \nabla^{-\delta_2} e^{-2\delta_1 \nabla^{-\delta_2}} \log(\nabla)}{(1 - e^{-2\delta_1 \nabla^{-\delta_2}})^2}, \end{aligned}$$

and

$$\begin{aligned} \frac{\partial \ell_n}{\partial \omega_k} &= -(\delta_2 + 1) \sum_{i=1}^n \partial(\log(\nabla)) / \partial \omega_k + \sum_{i=1}^n \partial(\log(\nabla')) / \partial \omega_k \\ &- 2 \sum_{i=1}^n \partial(\log(\delta_1 \nabla^{-\delta_2})) / \partial \omega_k + \sum_{i=1}^n \partial(\log(1 + e^{-2\delta_1 \nabla^{-\delta_2}})) / \partial \omega_k \\ &- 3 \sum_{i=1}^n \partial(\log(1 - e^{-2\delta_1 \nabla^{-\delta_2}})) / \partial \omega_k - \sum_{i=1}^n \partial \left(\left[\frac{e^{-2\delta_1 \nabla^{-\delta_2}}}{1 - e^{-2\delta_1 \nabla^{-\delta_2}}} \right] \right) / \partial \omega_k, \end{aligned}$$

where $\nabla = \frac{G(y;\omega)}{G^2(y;\omega)}$ and $\nabla' = \frac{g(y;\omega)}{G^2(y;\omega)}$. Therefore the estimates of the parameters are obtained by solving the non-linear systems of equation $\hat{\Delta} = \left(\frac{\partial \ell_n}{\partial \delta_1}, \frac{\partial \ell_n}{\partial \delta_2}, \frac{\partial \ell_n}{\partial \omega_k} \right)^T = \mathbf{0}$, using a numerical method such as Newton-Raphson technique [17, 18], via statistical softwares such as MATHEMATICA, MAPLE, Ox and R.

6. Simulation Study

In this section, a simulation exercise was conducted using the R programming language (stats4 package) to assess the consistency of the maximum likelihood estimators (MLEs). The OGR-RW-BXII distribution was used to generate 3000 samples with varying sizes ($n = 100, 200, 400, 800, 1000, 1600, 1800$ and 2000) through iterative simulations. For each of the 3000 replications, the MLEs were computed. The average bias (ABias) and root mean square error (RMSE) were calculated for the estimated parameters (refer to [14, 16] for the mathematical formulations of ABias and RMSE).

It is expected that as the sample size increases, both the ABias and RMSE will decrease toward zero, indicating good model performance. As shown in Tables 1 and 2, the ABias and RMSE for all parameter estimates consistently decline toward zero with increasing sample size. This demonstrates that the MLEs are consistent and produce reliable estimates for the model parameters, confirming the robustness of the model.

Table 1. OGR-RW-BXII Simulation Results 1

| parameter | Sample Size | (1.0, 1.0, 1.0, 1.0) | | | (1.0, 1.0, 1.5, 1.0) | | |
|------------|-------------|----------------------|--------|---------|----------------------|---------|--------|
| | | Mean | RMSE | ABias | Mean | RMSE | A.Bias |
| δ_1 | 100 | 1.7963 | 2.8691 | 0.7963 | 1.6712 | 3.6361 | 0.6712 |
| | 200 | 1.4372 | 1.8024 | 0.4372 | 1.4566 | 2.5401 | 0.4566 |
| | 400 | 1.2227 | 1.2856 | 0.2227 | 1.4837 | 2.4789 | 0.4837 |
| | 800 | 1.0722 | 0.4964 | 0.0722 | 1.2589 | 1.2322 | 0.2589 |
| | 1000 | 1.0573 | 0.4668 | 0.0573 | 1.2223 | 1.0633 | 0.2223 |
| | 1600 | 1.0253 | 0.2943 | 0.0253 | 1.1974 | 0.8863 | 0.1974 |
| | 1800 | 1.0361 | 0.2612 | 0.0361 | 1.1951 | 0.8478 | 0.1951 |
| | 2000 | 1.0299 | 0.2653 | 0.0299 | 1.2350 | 0.8681 | 0.2350 |
| δ_2 | 100 | 0.8811 | 0.3679 | -0.1189 | 0.8047 | -0.6712 | 0.1953 |
| | 200 | 0.9220 | 0.3131 | -0.0780 | 0.8411 | -0.6154 | 0.1589 |
| | 400 | 0.9705 | 0.2395 | -0.0295 | 0.9005 | -0.5531 | 0.0995 |
| | 800 | 0.9854 | 0.1493 | -0.0146 | 0.9184 | -0.5060 | 0.0816 |
| | 1000 | 0.9883 | 0.1331 | -0.0117 | 0.9421 | -0.4821 | 0.0579 |
| | 1600 | 0.9971 | 0.0899 | -0.0029 | 0.9630 | -0.4316 | 0.0370 |
| | 1800 | 0.9951 | 0.0656 | -0.0049 | 0.9496 | -0.4344 | 0.0504 |
| | 2000 | 0.9955 | 0.0777 | -0.0045 | 0.9672 | -0.4264 | 0.0328 |
| α | 100 | 1.3614 | 1.1162 | 0.3614 | 1.7869 | 1.7529 | 0.2869 |
| | 200 | 1.1819 | 0.7627 | 0.1819 | 1.5973 | 1.1795 | 0.0973 |
| | 400 | 1.0768 | 0.4658 | 0.0768 | 1.5642 | 0.9855 | 0.0642 |
| | 800 | 1.0310 | 0.2701 | 0.0310 | 1.4986 | -0.5367 | 0.0014 |
| | 1000 | 1.0172 | 0.2479 | 0.0172 | 1.4739 | -0.4672 | 0.0261 |
| | 1600 | 1.0086 | 0.1903 | 0.0086 | 1.4857 | -0.3765 | 0.0143 |
| | 1800 | 1.0188 | 0.1706 | 0.0188 | 1.4975 | -0.3615 | 0.0025 |
| | 2000 | 1.0142 | 0.1638 | 0.0142 | 1.5107 | 0.3539 | 0.0107 |
| β | 100 | 1.4381 | 1.5242 | 0.4381 | 3.2173 | 4.3280 | 2.2173 |
| | 200 | 1.3329 | 1.3022 | 0.3329 | 2.7617 | 3.5730 | 1.7617 |
| | 400 | 1.1425 | 0.7157 | 0.1425 | 2.2086 | 2.6175 | 1.2086 |
| | 800 | 1.0455 | 0.3623 | 0.0455 | 1.8826 | 1.9930 | 0.8826 |
| | 1000 | 1.0409 | 0.3122 | 0.0409 | 1.7821 | 1.8426 | 0.7821 |
| | 1600 | 1.0094 | 0.1643 | 0.0094 | 1.5556 | 1.4604 | 0.5556 |
| | 1800 | 0.9994 | 0.0927 | -0.0006 | 1.5595 | 1.4105 | 0.5595 |
| | 2000 | 1.0040 | 0.1327 | 0.0040 | 1.5074 | 1.3515 | 0.5074 |

Table 2. OGR-RW-BXII Simulation Results 2

| parameter | Sample Size | (1.0, 1.0, 1.2, 1.0) | | | (1.1, 1.1, 1.5, 1.1) | | |
|------------|-------------|----------------------|---------|--------|----------------------|---------|--------|
| | | Mean | RMSE | Bias | Mean | RMSE | A.Bias |
| δ_1 | 100 | 1.7641 | 3.6072 | 0.7641 | 1.7645 | 3.8650 | 0.6645 |
| | 200 | 1.4134 | 2.1130 | 0.4134 | 1.6397 | 3.3009 | 0.5397 |
| | 400 | 1.3403 | 1.7863 | 0.3403 | 1.4861 | 2.1106 | 0.3861 |
| | 800 | 1.1283 | 0.8190 | 0.1283 | 1.3035 | 1.1731 | 0.2035 |
| | 1000 | 1.1013 | 0.7398 | 0.1013 | 1.2955 | 1.1402 | 0.1955 |
| | 1600 | 1.0610 | 0.5193 | 0.0610 | 1.2596 | 0.9110 | 0.1596 |
| | 1800 | 1.0758 | 0.5158 | 0.0758 | 1.2742 | 0.8951 | 0.1742 |
| | 2000 | 1.0604 | 0.4850 | 0.0604 | 1.2892 | 0.8778 | 0.1892 |
| δ_2 | 100 | 0.8343 | -0.5182 | 0.1657 | 0.8910 | -0.7265 | 0.2090 |
| | 200 | 0.8870 | -0.4353 | 0.1130 | 0.9053 | -0.6524 | 0.1947 |
| | 400 | 0.9564 | -0.3752 | 0.0436 | 0.9910 | -0.5801 | 0.1090 |
| | 800 | 0.9871 | -0.2943 | 0.0129 | 1.0033 | -0.5247 | 0.0967 |
| | 1000 | 0.9811 | -0.2868 | 0.0189 | 1.0377 | -0.5043 | 0.0623 |
| | 1600 | 0.9958 | -0.2161 | 0.0042 | 1.0556 | -0.4461 | 0.0444 |
| | 1800 | 0.9901 | -0.2229 | 0.0099 | 1.0350 | -0.4307 | 0.0650 |
| | 2000 | 0.9947 | -0.2047 | 0.0053 | 1.0539 | -0.4291 | 0.0461 |
| α | 100 | 1.5397 | 1.4386 | 0.3397 | 1.7967 | 1.7967 | 0.2967 |
| | 200 | 1.3428 | 0.8876 | 0.1428 | 1.6365 | 1.3629 | 0.1365 |
| | 400 | 1.2814 | 0.6875 | 0.0814 | 1.5326 | 0.8775 | 0.0326 |
| | 800 | 1.2194 | 0.3536 | 0.0194 | 1.4795 | -0.4979 | 0.0205 |
| | 1000 | 1.2055 | 0.3209 | 0.0055 | 1.4651 | -0.4630 | 0.0349 |
| | 1600 | 1.2046 | 0.2473 | 0.0046 | 1.4745 | -0.3710 | 0.0255 |
| | 1800 | 1.2168 | 0.2280 | 0.0168 | 1.4935 | -0.3534 | 0.0065 |
| | 2000 | 1.2095 | 0.2184 | 0.0095 | 1.4940 | -0.3400 | 0.0060 |
| β | 100 | 1.9868 | 2.3237 | 0.9868 | 3.4431 | 4.6703 | 2.3431 |
| | 200 | 1.6594 | 1.6981 | 0.6594 | 2.8542 | 3.5304 | 1.7542 |
| | 400 | 1.3901 | 1.2132 | 0.3901 | 2.2719 | 2.5523 | 1.1719 |
| | 800 | 1.1884 | 0.7732 | 0.1884 | 1.9650 | 1.9745 | 0.8650 |
| | 1000 | 1.1968 | 0.7513 | 0.1968 | 1.8443 | 1.7809 | 0.7443 |
| | 1600 | 1.0879 | 0.4855 | 0.0879 | 1.6265 | 1.4069 | 0.5265 |
| | 1800 | 1.0964 | 0.5083 | 0.0964 | 1.6240 | 1.3590 | 0.5240 |
| | 2000 | 1.0772 | 0.4546 | 0.0772 | 1.5792 | 1.3033 | 0.4792 |

7. Application

This section demonstrates the utility of the OGR-RW-G family by applying the OGR-RW-BXII distribution to fit three distinct datasets. The model's performance is assessed using various goodness-of-fit (GoF) statistics, including $-2 \log$ -likelihood ($-2 \log L$), Akaike Information Criterion (AIC), Consistent Akaike Information Criterion (AICC), Bayesian Information Criterion (BIC), Cramér-von Mises (W^*), Anderson-Darling (A^*), and the Kolmogorov-Smirnov (K-S) statistic along with its p-value. Among the models considered, the one with the lowest values for these statistics and the highest K-S p-value is deemed the most optimal.

The OGR-RW-BXII distribution is compared to its nested models as well as five competing non-nested five-parameter models: the Beta Generalized Lindley (BGL) distribution [19], Generalized Gompertz-Poisson (GGP) distribution [8], Beta Odd Lindley-Exponential (BOLE) distribution [5], Weibull Lomax (WL) distribution [21], and Marshall-Olkin-Gompertz-Weibull (MO-Gom-W) distribution [6].

Parameter estimates, along with their standard errors (in parentheses), for the selected datasets are provided in Tables 3 and 7, while the corresponding GoF statistics are summarized in Tables 4 and 6. Figures 5 and 9 illustrate the fitted density, probability plots, and Kaplan-Meier (KM) curves for the OGR-RW-BXII model. Additionally, Figures 6 and 10 present the empirical cumulative distribution function (ECDF), estimated hazard rate function (HRF), and total-time-on-test (TTT) plots for the OGR-RW-BXII model.

These analyses highlight the flexibility and effectiveness of the OGR-RW-BXII model in accurately fitting diverse datasets.

7.1. Repair Lifetimes of an Airborne Transceiver

The dataset represents the duration of active repairs measured in hours for an airborne communication transceiver. These data points are sourced from the studies conducted by Raj S. Chhikara and J Leroy Folks [4] and Victor Leiva et.al. [9]. The variance-covariance matrix for OGR-RW-BXII model on Repair lifetimes of an airborne transceiver dataset is given by

$$\begin{bmatrix} 1.4060 \times 10^{-13} & 3.2053 \times 10^{-16} & -8.8983 \times 10^{-13} & -1.2463 \times 10^{-17} \\ 3.2053 \times 10^{-16} & 7.3069 \times 10^{-19} & -2.0284 \times 10^{-15} & -2.8412 \times 10^{-20} \\ -8.8982 \times 10^{-13} & -2.0285 \times 10^{-15} & 5.6312 \times 10^{-12} & 7.8875 \times 10^{-17} \\ -1.2463 \times 10^{-17} & -2.8412 \times 10^{-20} & 7.8875 \times 10^{-17} & 1.1048 \times 10^{-21} \end{bmatrix}$$

and the 95% confidence intervals for the model parameters are given by $\delta_1 \in [2.2552 \times 10^{-4} \pm 7.3495 \times 10^{-7}]$, $\delta_2 \in [0.7492 \pm 1.6754 \times 10^{-6}]$, $\alpha \in [2.6700 \times 10^{-5} \pm 4.6511 \times 10^{-6}]$ and $\beta \in [1.0424 \pm 6.5147 \times 10^{-11}]$. Table 4 shows that the OGR-RW-BXII model provides the overall best fit on the repair lifetimes of an airborne transceiver dataset as compared to nested and non-nested models presented in the table.

Figures 5 and 6 show that the OGR-RW-BXII model can accommodate the extremely tailed data sets, KM and ECDF curves are close to the empirical data exhibiting better performance of our novel model. Furthermore, the fitted hrf exhibits an upside-down bathtub shape which is supported by the TTT plot produced by the dataset.

7.2. Growth Hormone Dataset

The second dataset comprises the estimated duration from the administration of growth hormone medication until the children reached the desired age [13].

The variance-covariance matrix for OGR-RW-BXII model on growth hormone data set set is given by

$$\begin{bmatrix} 1.406 \times 10^{-13} & 3.205 \times 10^{-16} & -8.898 \times 10^{-13} & -1.246 \times 10^{-17} \\ 3.205 \times 10^{-16} & 7.307 \times 10^{-19} & -2.028 \times 10^{-15} & -2.841 \times 10^{-20} \\ -8.898 \times 10^{-13} & -2.028 \times 10^{-15} & 5.631 \times 10^{-12} & 7.888 \times 10^{-17} \\ -1.246 \times 10^{-17} & -2.841 \times 10^{-20} & 7.888 \times 10^{-17} & 1.105 \times 10^{-21} \end{bmatrix}$$

Table 3. Fitted models parameter estimates for the repair lifetimes of an airborne transceiver dataset

| Model | Estimates and standard errors | | | |
|---|--|--|--|--|
| OGR-RW-BXII | δ_1 | δ_2 | α | β |
| | 2.2552×10^{-4} (3.7497×10^{-7}) | 0.7492 (8.5481×10^{-10}) | 2.6700×10^{-5} (2.3730×10^{-6}) | 1.0424 (3.3238×10^{-11}) |
| OGR-RW-BXII($1, \delta_2, \alpha, \beta$) | - | 14.7030 (4.1329×10^{-6}) | 1.0104 (0.0048) | 2.8863 (0.0029) |
| OGR-RW-BXII($\delta_1, 1, \alpha, \beta$) | 3.0031×10^{-5} (1.5652×10^{-6}) | - | 5.3642×10^{-5} (8.7628×10^{-7}) | 0.7338 (3.1085×10^{-11}) |
| BGL | α | θ | a | b |
| | 1.1741×10^{-1} (0.0010) | 1.3021×10^{-7} (1.8480×10^{-6}) | 0.0408 (6.0198×10^{-3}) | 8.0996 (2.9912×10^{-5}) |
| GGP | θ | α | β | γ |
| | 1.5838×10^{-4} (1.0223) | 0.8065 (0.2288) | 0.2266 (0.0668) | 3.9440×10^{-9} (0.0180) |
| BOLE | λ | θ | a | b |
| | 80.209 (1.2470×10^{-11}) | 8.0481×10^{-5} (1.2289×10^{-5}) | 0.9317 (1.0488×10^{-9}) | 40.511 (2.4410×10^{-11}) |
| WL | a | b | α | β |
| | 2.0041×10^2 (0.0017) | 1.1500 (0.2283) | 0.1738 (0.0086) | 3.7634 (3.6149) |
| MO-GOm-W | δ | θ | λ | γ |
| | 0.0273 (0.0747) | 0.0122 (0.0342) | 1.1551 (0.1734) | 0.0403 (0.0632) |
| G | | | a | b |
| | | | 0.2585 (0.0615) | 0.9323 (0.1701) |
| W | | | | λ |
| | | | | 0.5649 (0.0500) |

Table 4. Fitted models GoF statistics for the repair lifetimes of an airborne transceiver dataset

| Model | GoF Statistics | | | | | | | |
|---|----------------|----------|----------|----------|--------|--------|--------|-----------------------|
| | $-2 \log L$ | AIC | $AICC$ | BIC | W^* | A^* | K-S | p-value |
| OGR-RW-BXII | 199.3641 | 207.3641 | 208.3397 | 214.6786 | 0.0460 | 0.2841 | 0.0928 | 0.8235 |
| OGR-RW-BXII($1, \delta_2, \alpha, \beta$) | 204.769 | 471.4273 | 471.9988 | 476.9133 | 0.0527 | 0.3351 | 0.7875 | 2.2×10^{-16} |
| OGR-RW-BXII($\delta_1, 1, \alpha, \beta$) | 199.8261 | 205.8261 | 206.3975 | 211.3120 | 0.0501 | 0.3234 | 0.1043 | 0.6989 |
| BGL | 539.3437 | 547.6217 | 548.5973 | 554.9363 | 0.0742 | 0.4838 | 0.9561 | 2.2×10^{-16} |
| GGP | 210.8166 | 218.8167 | 219.7923 | 226.1313 | 0.1441 | 1.0013 | 0.1334 | 0.3856 |
| BOLE | 209.8717 | 217.8717 | 218.8473 | 225.1862 | 0.1434 | 0.9950 | 0.1455 | 0.2842 |
| WL | 204.2995 | 212.2995 | 213.2751 | 219.6140 | 0.0803 | 0.5158 | 0.1157 | 0.5692 |
| MO-GOm-W | 204.0075 | 212.0075 | 212.9831 | 219.3221 | 0.0683 | 0.4692 | 0.1251 | 0.4675 |
| G | 209.8619 | 213.8619 | 214.1410 | 217.5192 | 0.1069 | 0.6729 | 0.1603 | 0.1877 |
| W | 241.9077 | 243.9077 | 243.9986 | 245.7364 | 0.0978 | 0.6719 | 0.4479 | 1.9×10^{-8} |

and the 95% confidence intervals for the model parameters are given by $\delta_1 \in [2.151 \times 10^{-4} \pm 7.349 \times 10^{-07}]$, $\delta_2 \in [1.569 \pm 1.675 \times 10^{-09}]$, $\alpha \in [3.219 \times 10^{-4} \pm 4.651 \times 10^{-06}]$ and $\beta \in [13.316 \pm 6.515 \times 10^{-11}]$. Table 6 shows that the OGR-RW-BXII model provides the overall best fit on the




Figure 5. Fitted density, probability plot and KM survival plots of the OGR-RW-BXII model for repair lifetimes of an airborne transceiver data




Figure 6. ECDF, estimated hazard rate plot and TTT of the OGR-RW-BXII model for repair lifetimes of an airborne transceiver data

growth hormone dataset as compared to the nested and non-nested models under consideration. Figures 9 and 10




Figure 7. Fitted density, probability plots and KM survival plots of the OGR-RW-BXII model for the growth hormone data

that the OGR-RW-BXII model can accommodate the extremely tailed data sets, the KM and ECDF are aligning closely with the model thus exhibiting better performance for our model. Additionally, the estimated hrf exhibits the TTT plot produced by the dataset.

Table 5. Fitted models parameter estimates for the growth hormone dataset

| Model | Estimates and standard errors | | | |
|---|-------------------------------|---------------------------|---------------------------|---------------------------|
| OGR-RW-BXII | δ_1 | δ_2 | α | β |
| | 2.1510×10^{-4} | 1.5690 | 3.2190×10^{-4} | 13.3160 |
| | (6.4410×10^{-4}) | (2.0940×10^{-6}) | (6.1250×10^{-4}) | (6.9100×10^{-8}) |
| OGR-RW-BXII($1, \delta_2, \alpha, \beta$) | - | 0.9760 | 0.0460 | 14.0380 |
| | | (0.1450) | (0.0040) | (1.4080×10^{-5}) |
| OGR-RW-BXII($\delta_1, 1, \alpha, \beta$) | 0.0670 | - | 0.1110 | 0.4910 |
| | (0.1070) | | (0.1720) | (0.0850) |
| BG-L | α | θ | a | b |
| | 0.1151 | 1.6105×10^{-7} | 0.0391 | 8.0101 |
| | (9.6102×10^{-3}) | (1.9382×10^{-6}) | (6.5904×10^{-3}) | (3.4570×10^{-5}) |
| GG-P | θ | α | β | γ |
| | 5.2833×10^{-9} | 1.4484 | 0.1400 | 0.1300 |
| | (0.0101) | (0.4664) | (0.0631) | (0.0590) |
| BOL-E | λ | θ | a | b |
| | 24.2867 | 0.0025 | 4.1427 | 12.8370 |
| | (0.004) | (4.988×10^{-4}) | (0.9550) | (0.0024) |
| W-L | a | b | α | β |
| | 546.4500 | 4.4581 | 0.0664 | 0.8486 |
| | (6.6900×10^{-5}) | (3.2384) | (0.0166) | (2.2971) |
| MO-GOm-W | δ | θ | λ | γ |
| | 0.0370 | 2.1544×10^{-5} | 0.3410 | 5.3427 |
| | (0.0107) | (5.9010×10^{-6}) | (0.0186) | (7.1244×10^{-4}) |
| G | | | a | b |
| | | | 0.7861 | 4.1711 |
| | | (0.1923) | (0.9599) | |
| W | | | | λ |
| | | | | 0.4836 |
| | | | | (0.0513) |

Table 6. Fitted models GoF statistics for the Growth hormone data set

| Model | Statistics | | | | | | | |
|---|-------------|----------|----------|----------|--------|--------|--------|------------------------|
| | $-2 \log L$ | AIC | CAIC | BIC | W^* | A^* | K-S | p-value |
| OGR-RW-BXII | 154.4357 | 162.4358 | 163.7691 | 168.6572 | 0.0376 | 0.2584 | 0.0822 | 0.9721 |
| OGR-RW-BXII($1, \delta_2, \alpha, \beta$) | 156.2151 | 386.5832 | 387.3574 | 391.2493 | 0.0329 | 0.2332 | 0.8055 | 2.2×10^{-16} |
| OGR-RW-BXII($\delta_1, 1, \alpha, \beta$) | 209.3691 | 215.3691 | 216.1433 | 220.0351 | 0.0610 | 0.4173 | 0.4160 | 1.095×10^{-5} |
| BG-L | 471.1036 | 479.1123 | 480.4456 | 485.3337 | 0.0627 | 0.4285 | 0.9582 | 2.2×10^{-16} |
| GG-P | 169.0884 | 177.0885 | 178.4218 | 183.3099 | 0.1986 | 1.2386 | 0.1702 | 0.2625 |
| BOL-E | 160.2746 | 168.2746 | 169.6079 | 174.496 | 0.1019 | 0.6609 | 0.1251 | 0.6438 |
| W-L | 161.0264 | 169.0264 | 170.3597 | 175.2478 | 0.1098 | 0.7068 | 0.1244 | 0.6508 |
| MO-GOm-W | 161.5606 | 169.5607 | 170.8940 | 175.782 | 0.1134 | 0.7346 | 0.1037 | 0.846 |
| G | 160.2165 | 164.2165 | 164.5915 | 167.3272 | 0.0816 | 0.5885 | 0.8671 | 2.2×10^{-16} |
| W | 258.6514 | 260.6514 | 260.7727 | 262.2068 | 0.0771 | 0.5138 | 0.7650 | 2.2×10^{-16} |




Figure 8. ECDF, estimated hazard rate plot and TTT of the OGR-RW-BXII model for the growth hormone data

7.3. Survival Dataset of Patients Suffering from Acute Myelogeneous Leukaemia

The third dataset comprises of patients suffering from acute myelogeneous leukaemia. The data, that can also be found at library SMIR of the R program (<http://cran.r-project.org>). The variance-covariance matrix for OGR-RW-BXII model on myelogeneous leukaemia dataset set is given by

$$\begin{bmatrix} 8.7490 \times 10^{-2} & -4.7800 \times 10^{-5} & 1.5970 \times 10^{-3} & 0.8730 \\ -4.7800 \times 10^{-5} & 5.6330 \times 10^{-4} & -2.3720 \times 10^{-4} & -0.1958 \\ 1.5970 \times 10^{-3} & -2.3720 \times 10^{-4} & 0.0004 & -0.1737 \\ 0.8732 & -0.1958 & -0.1737 & 310.0262 \end{bmatrix}$$

and the 95% confidence intervals for the model parameters are given by $\delta_1 \in [1.5252 \times 10^{-4} \pm 0.5797]$, $\delta_2 \in [0.0817 \pm 0.0465]$, $\alpha \in [0.0687 \times 10^{-4} \pm 0.0421]$ and $\beta \in [57.6924 \pm 34.5108]$. Table 8 shows that the OGR-RW-BXII model provides the overall best fit on the myelogeneous leukaemia dataset as compared to the nested and non-nested models under consideration. Figures 9 and 10 that the




Figure 9. Fitted density, probability plots and KM survival plots of the OGR-RW-BXII model for the myelogeneous leukaemia data

OGR-RW-BXII model can accommodate the extremely tailed data sets, the KM and ECDF are aligning closely with the model thus exhibiting better performance for our model. Additionally, the estimated hrf exhibits the TTT plot produced by the dataset.

Table 7. Fitted models parameter estimates for the myelogeneous leukaemia dataset

| Model | Estimates and standard errors | | | |
|---|-------------------------------|----------------------------|-----------------------------|-----------------------------|
| OGR-RW-BXII | δ_1 | δ_2 | α | β |
| | 1.5252 | 0.0817 | 0.0687 | 57.6924 |
| | (0.2958) | (0.0237) | (0.0215) | (17.6076) |
| OGR-RW-BXII($1, \delta_2, \alpha, \beta$) | - | 0.4183 | 0.2355 | 2.4818 |
| | | (0.1594) | (0.1780) | (2.5172) |
| OGR-RW-BXII($\delta_1, 1, \alpha, \beta$) | 1.0610 | - | 0.7025 | 0.4171 |
| | (1.9050) | | (1.0600) | (0.2960) |
| BG-L | α | θ | a | b |
| | 0.1034 | 1.6105×10^{-8} | 0.0331 | 8.0997 |
| | (8.378×10^{-3}) | (1.409×10^{-6}) | (5.766×10^{-3}) | (3.129×10^{-5}) |
| GG-P | θ | α | β | γ |
| | 7.142×10^{-5} | 0.5478 | 0.0124 | 0.0056 |
| | (0.1274) | (0. 1548) | (0.0077) | (0.0073) |
| BOL-E | λ | θ | a | b |
| | 1.5185 | 0.0028 | 0.6204 | 4.9850 |
| | (2.2114) | (0.0045) | (0.1732) | (0.4522) |
| W-L | a | b | α | β |
| | 1.2047 | 0.6017 | 1.510×10^3 | 1.156×10^3 |
| | (0.2141) | (0.0805) | (1.6227×10^{-4}) | (2.1186×10^{-6}) |
| MO-GOm-W | δ | θ | λ | γ |
| | 0.1715 | 0.0106 | 0.8313 | 0.0263 |
| | (0.2648) | (0.0181) | (0.2039) | (0.0401) |
| G | | | a | b |
| | | | 0.0168 | 0.6878 |
| | | (0.0050) | (0.1440) | |
| W | | | λ | |
| | | | 0.2393 | |
| | | | (0.0259) | |

Table 8. Fitted models GoF statistics for the myelogeneous leukaemia dataset

| Model | Statistics | | | | | | | |
|---|-------------|----------|----------|----------|--------|--------|--------|------------------------|
| | $-2 \log L$ | AIC | $CAIC$ | BIC | W^* | A^* | K-S | p-value |
| OGR-RW-BXII | 298.9213 | 306.9213 | 308.3499 | 312.9074 | 0.0705 | 0.4767 | 0.1267 | 0.6644 |
| OGR-RW-BXII($1, \delta_2, \alpha, \beta$) | 304.6768 | 339.974 | 340.8016 | 344.4635 | 0.1097 | 0.7076 | 0.4068 | 3.62×10^{-5} |
| OGR-RW-BXII($\delta_1, 1, \alpha, \beta$) | 305.3852 | 311.3852 | 312.2128 | 315.8747 | 0.0862 | 0.5710 | 0.1440 | 0.5008 |
| BG-L | 549.7148 | 557.8677 | 559.2963 | 563.8538 | 0.1276 | 0.8035 | 0.9644 | 2.2×10^{-16} |
| GG-P | 307.1409 | 315.1409 | 316.5694 | 321.1269 | 0.1030 | 0.7129 | 0.1172 | 0.7556 |
| BOL-E | 307.0134 | 315.0134 | 316.4420 | 320.9995 | 0.0987 | 0.6849 | 0.1369 | 0.5664 |
| W-L | 306.4082 | 376.8067 | 376.9357 | 378.3032 | 0.1001 | 0.6548 | 0.6368 | 4.78×10^{-12} |
| MO-GOm-W | 305.3181 | 313.3181 | 314.7467 | 319.3041 | 0.0857 | 0.5889 | 0.1312 | 0.6206 |
| G | 307.3473 | 311.3473 | 311.7473 | 314.3403 | 0.1284 | 0.8533 | 0.2695 | 0.0166 |
| W | 374.8067 | 260.6514 | 260.7727 | 262.2068 | 0.0771 | 0.5138 | 0.7650 | 2.2×10^{-16} |




Figure 10. ECDF, estimated hazard rate plot and TTT of the OGR-RW-BXII model for the myelogenous leukaemia data

8. Concluding Remarks

In this study, we introduced the Odd Generalized Rayleigh Reciprocal Weibull (OGR-RW-G) family of probability distributions, representing a novel and versatile class of probability models. The statistical properties of this newly developed family were thoroughly analyzed, and key characteristics were derived. The maximum likelihood estimation (MLE) method was employed to estimate the model parameters, and a Monte Carlo simulation study was conducted to evaluate the performance of the MLEs.

Additionally, the empirical performance of the OGR-RW-BXII distribution, a member of the OGR-RW-G family, was assessed using real-world datasets. The results demonstrate that the OGR-RW-BXII distribution outperforms its nested distributions as well as other non-nested competing distributions under consideration. These findings underscore the flexibility and effectiveness of the OGR-RW-G family in modeling complex datasets and provide a foundation for its application in various fields requiring robust statistical modeling.

REFERENCES

1. Konstantinos Adamidis and S Loukas. *A lifetime distribution with decreasing failure rate*. *Statistics & Probability Letters*, 39(1):35–42, 1998.
2. Ali Al-Shomrani, Osama Arif, A Shawky, Saman Hanif, and Muhammad Qaiser Shahbaz. *Toppâ€“Leone Family of Distributions: Some Properties and Application*. *Pakistan Journal of Statistics and Operation Research*, pages 443–451, 2016.
3. Ayman Alzaatreh, Carl Lee, and Felix Famoye. *A new method for generating families of continuous distributions*. *Metron*, 71(1):63–79, 2013.
4. Raj S Chhikara and J Leroy Folks. *The inverse gaussian distribution as a lifetime model*. *Technometrics*, 19(4):461–468, 1977.
5. F Chipepa, B Oluyede, B Makubate, and AF Fagbamigbe. *The beta odd lindley-g family of distributions with applications*. *Journal of Probability and Statistical Science*, 17(1):51–83, 2019.
6. Fastel Chipepa and Broderick Oluyede. *The marshall-olkin-gompertz-g family of distributions: properties and applications*. *J. Nonlinear Sci. Appl*, 14(4):257–260, 2021.
7. Gauss M Cordeiro, Edwin MM Ortega, and Daniel CC da Cunha. *The exponentiated generalized class of distributions*. *Journal of data science*, 11(1):1–27, 2013.
8. Awad El-Gohary, Ahmad Alshamrani, and Adel Naif Al-Otaibi. *The generalized gompertz distribution*. *Applied mathematical modelling*, 37(1-2):13–24, 2013.
9. Victor Leiva, Michelli Karinne Barros da Silva, and Gilberto Alvarenga Paula. *Generalized birnbaum-saunders models using r*. 2009.
10. Boikanyo Makubate and Regent Retrospect Musekwa. *A novel technique for generating families of continuous distributions*. *Statistics, Optimization & Information Computing*, 12(5):1231–1248, 2024.
11. Boikanyo Makubate, Broderick Oluyede, and Morongwa Gabanakgosi. *A New Lindley-Burr XII Distribution: Model, Properties and Applications*. *International Journal of Statistics and Probability*, 10(4):33–51, 2021.
12. Albert W Marshall and Ingram Olkin. *A new method for adding a parameter to a family of distributions with application to the exponential and Weibull families*. *Biometrika*, 84(3):641–652, 1997.
13. Thatayaone Moakofi, Broderick Oluyede, and Morongwa Gabanakgosi. *The topp-leone odd burr iii-g family of distributions: Model, properties and applications*. *Statistics, Optimization & Information Computing*, 10(1):236–262, 2022.
14. Regent Retrospect Musekwa, Lesego Gabaitiri, and Boikanyo Makubate. *Application of the marshall-olkin-weibull logarithmic distribution to complete and censored data*. *Heliyon*, 10(14), 2024.

15. Regent Retrospect Musekwa, Gabaitiri Lesego, and Boikanyo Makubate. A new technique to generate families of continuous distributions. *Revista Colombiana de Estadística*, 47(2):1–24, 2024.
16. Regent Retrospect Musekwa and Boikanyo Makubate. Statistical analysis of saudi arabia and uk covid-19 data using a new generalized distribution. *Scientific African*, 22:e01958, 2023.
17. Regent Retrospect Musekwa and Boikanyo Makubate. A flexible generalized x lindley distribution with application to engineering. *Scientific African*, page e02192, 2024.
18. Violet Zivai Nyamajiwa, Regent Retrospect Musekwa, and Boikanyo Makubate. Application of the new extended topp-leone distribution to complete and censored data. *Revista Colombiana de Estadística*, 47(1):37–65, 2024.
19. Broderick O Oluyede and Tiantian Yang. A new class of generalized lindley distributions with applications. *Journal of Statistical Computation and Simulation*, 85(10):2072–2100, 2015.
20. Miroslav M Ristić and Narayanaswamy Balakrishnan. *The gamma-exponentiated exponential distribution*. *Journal of statistical computation and simulation*, 82(8):1191–1206, 2012.
21. Muhammad H Tahir, Gauss M Cordeiro, M Mansoor, and Muhammad Zubair. The weibull-lomax distribution: properties and applications. *Hacettepe Journal of Mathematics and Statistics*, 44(2):455–474, 2015.
22. Chester W Topp and Fred C Leone. *A family of J-shaped frequency functions*. *Journal of the American Statistical Association*, 50(269):209–219, 1955.
23. Konstantinos Zografos and Narayanaswamy Balakrishnan. *On families of beta-and generalized gamma-generated distributions and associated inference*. *Statistical methodology*, 6(4):344–362, 2009.